

Asymptotic finite-time ruin probabilities for a class of path-dependent heavy-tailed claim amounts using Poisson spacings

Romain Biard^{a,*†}, Claude Lefèvre^b, Stéphane Loisel^a and Haikady N. Nagaraja^c

In the compound Poisson risk model, several strong hypotheses may be found too restrictive to describe accurately the evolution of the reserves of an insurance company. This is especially true for a company that faces natural disaster risks like earthquake or flooding. For such risks, claim amounts are often inter-dependent and they may also depend on the history of the natural phenomenon. The present paper is concerned with a situation of this kind, where each claim amount depends on the previous claim inter-arrival time, or on past claim inter-arrival times in a more complex way. Our main purpose is to evaluate, for large initial reserves, the asymptotic finite-time ruin probabilities of the company when the claim sizes have a heavy-tailed distribution. The approach is based more particularly on the analysis of spacings in a conditioned Poisson process. Copyright © 2010 John Wiley & Sons, Ltd.

Keywords: risk process; finite-time ruin probabilities; asymptotic approximation for large initial reserves; path-dependent claim amount; heavy-tailed claim amount; Poisson spacings

1. Introduction

The compound Poisson risk model is the central model proposed in the insurance theory. An abundant literature has been devoted to its analysis and applications. The reader is referred e.g. to the comprehensive books by Rolski *et al.* [1], Asmussen [2] and Goovaerts *et al.* [3]. It is well recognized, however, that the simplifying hypotheses at the basis of the model may be too restrictive for certain insurance coverages, especially with rare but extreme risks like earthquake or flooding. The present paper is concerned with such situations.

1.1. Framework and motivations

Let us begin by recalling the model in its standard version. The reserves of the company, $\{R(t), t \geq 0\}$, are given by

$$R(t) = u + ct - S(t), \quad t \geq 0, \quad (1)$$

where $u \geq 0$ is the amount of initial reserves, $c > 0$ is the premium income rate and $S(t)$ is the total claim amount up to time t . $\{S(t), t \geq 0\}$ is a compound Poisson process, i.e.

$$S(t) = \sum_{i=1}^{N(t)} X_i, \quad t \geq 0, \quad (2)$$

where $\{N(t), t \geq 0\}$ is a Poisson process, with parameter λ , which counts the claim occurrences until time t , and $\{X_i, i = 1, 2, \dots\}$ are the successive claim amounts which are represented by non-negative independent identically distributed random variables. Let $\{V_i, i = 1, 2, \dots\}$ be the inter-arrival times of successive claims. By construction, these random variables are

^a Université de Lyon, Université Lyon 1, Institut de Science Financière et d'Assurances, 50 Avenue Tony Garnier, F-69007 Lyon, France

^b Université Libre de Bruxelles, Département de Mathématique, Campus de la Plaine C.P. 210, B-1050 Bruxelles, Belgique

^c Ohio State University, Department of Statistics, 440H Cockins Hall, 1958 Neil Avenue, Columbus OH 43210-1247, U.S.A.

*Correspondence to: Romain Biard, Université de Lyon, Université Lyon 1, Institut de Science Financière et d'Assurances, 50 Avenue Tony Garnier, F-69007 Lyon, France.

†E-mail: romain.biard@univ-lyon1.fr

independent with a common exponential distribution. Moreover, the inter-occurrence times are assumed to be independent of the claim amounts.

A statistics of great interest is the probability of (non-)ruin over any fixed horizon of finite length. Denote by $\psi(u, t)$ the probability of ruin before time t , $t \geq 0$, for initial reserves u :

$$\psi(u, t) = P(\exists s \in [0, t], R(s) < 0 \mid R(0) = u), \quad u, t \geq 0. \quad (3)$$

Different methods have been proposed to evaluate $\psi(u, t)$; see, e.g. Lefèvre and Loisel [4].

Recently, much research has been devoted to the evaluation of ruin probabilities, over finite or infinite horizon, when some independence and stationarity assumptions of the model are relaxed. A number of references will be mentioned later in the section (the list being non-exhaustive, of course).

In practice, the independence assumptions on, and between, the sequences $\{X_i\}$ and $\{V_i\}$ may be too unrealistic. Typically, this arises in the case of natural disasters like earthquake (or flooding). The occurrence of an earthquake often increases the probability of by-claims in a near future. If the last earthquake occurred a long time ago, the next earthquake is likely to be more severe. If two earthquakes occur in a short time span, the second one may cause unusual damages like flooding.

This question has already been raised and discussed in the literature. Albrecher and Boxma [5] consider that the time between two claim occurrences depends on the previous claim amount. Exact expressions for the Laplace transform of the (ultimate) survival probability are derived. In Albrecher and Teugels [6], the claim inter-arrival time and the subsequent claim size are dependent on an arbitrary copula structure. Asymptotic results for both the finite and infinite-time ruin probabilities are then derived. Boudreault *et al.* [7] consider a particular form of dependence among the claim inter-arrival time and the subsequent claim size: if the current claim inter-arrival time exceeds a certain threshold, the distribution of the next claim is modified. The defective renewal equation satisfied by the expected Gerber–Shiu discounted penalty function is then obtained. In Meng *et al.* [8], the time between two claim occurrences determines the distribution of the next claim. Some exact and approximation results are derived for the survival probability. Ambagaspitiya [9] determines the ruin probability for two forms of dependence between claim size and occurrence in the Sparre Andersen model. See also the references given in these papers.

The present paper discusses several scenarios of dependence between claim amounts and claim inter-arrival time. Such scenarios are motivated by earthquake or flooding-type risks. Our purpose is to provide approximations to the finite-time ruin probabilities, for large initial reserves, when the claim amounts have heavy-tailed distributions. This work can also be seen as a sequel to a recent paper by Biard *et al.* [10]. Here, the claim amount distributions are allowed to depend, to some extent, on the history of the claim arrival process.

1.2. Basic assumptions and implications

As in the classical model, claims occur according to a Poisson process $\{N(t), t \geq 0\}$. Let $U_i = \sum_{j=1}^i V_j$, $i \geq 1$, be the claim arrival times.

Let us turn to the claim amounts $\{X_i, i \geq 1\}$. First, the X_i is allowed to be one of two different types of random variables Y_i or Z_i depending on the behavior of past inter-arrival times V_j , $j \leq i$. Three different models that describe this dependence will be examined.

Second, all the severities Y_i have the same distribution function (d.f.) F , and each vector $\mathbf{Y}^{(j)} = (Y_1, \dots, Y_j)$, $j \geq 1$, is of multivariate regular variation of index $-\alpha$ with $\alpha > 0$, i.e. there exists a $\theta \in \mathbf{S}^{j-1}$, where \mathbf{S}^{j-1} is the unit sphere with respect to a norm $|\bullet|$, such that

$$\frac{P(|\mathbf{Y}^{(j)}| > tx, \mathbf{Y}^{(j)} / |\mathbf{Y}^{(j)}| \in \bullet)}{P(|\mathbf{Y}^{(j)}| > x)} \xrightarrow{v} t^{-\alpha} P_{\mathbf{S}^{j-1}}(\theta \in \bullet),$$

where \xrightarrow{v} denotes vague convergence on \mathbf{S}^{j-1} (see e.g. Resnick [11] and Basrak *et al.* [12]). Analogously, the amounts Z_i too have a common distribution function G , and each vector $\mathbf{Z}^{(j)} = (Z_1, \dots, Z_j)$, $j \geq 1$, is again multivariate regularly varying but of index $-\beta$ with $\beta > 0$. The sequences $\{Y_i, i \geq 1\}$, $\{Z_i, i \geq 1\}$ and $\{V_i, i \geq 1\}$ are independent of each other.

A key tool in the analysis will be the following well-known result (see e.g. Barbe *et al.* [13]). If (Y_1, \dots, Y_j) is regularly varying of index $-\alpha$ with common marginal d.f. F , then the right tail of the partial sum $S_j = Y_1 + \dots + Y_j$ can be approximated as

$$P(Y_1 + \dots + Y_j > x) \sim q_{j,\alpha} \overline{F}(x) \quad \text{for large } x, \quad (4)$$

where \sim means that the ratio tends to 1 as $x \rightarrow \infty$. The parameter $q_{j,\alpha}$ in (4) depends on j , α and on the dependence structure of the vector (Y_1, \dots, Y_j) .

Table I provides examples of $q_{j,\alpha}$ for some classical multivariate copulas (see Nelsen [14] for a nice introduction to copulas). The independent copula corresponds to the special case where the Y_i , $1 \leq i \leq j$, are independent. The Fréchet upper

Table I. Values of $q_{j,\alpha}$ in (4) for different copulas.	
Special copulas	$q_{j,\alpha}$
Independent	j
Fréchet upper bound	j^α
Non-degenerate Gaussian	j
BLL	$\sum_{i=0}^j \binom{j}{i} p^i (1-p)^{j-i} (j-i+i^\alpha)$

bound represents the case where the Y_i 's are comonotonic. The non-degenerate Gaussian copula is the case where the dependence between the Y_i 's is drawn from a multivariate Gaussian distribution with a correlation matrix with coefficients strictly other than ± 1 outside of the diagonal. A copula introduced in [10], denoted BLL in the sequel, is built as follows:

$$Y_i = I_i W_0 + (1 - I_i) W_i, \quad 1 \leq i \leq j, \tag{5}$$

where $W_i, 0 \leq i \leq j$, are i.i.d. random variables, and $I_i, 1 \leq i \leq j$, are i.i.d. Bernoulli random variables with parameter p , these two sequences being mutually independent.

Let us mention that the study of the sum of dependent random variables has received much attention in actuarial sciences. See e.g. Wüthrich [15], Alink *et al.* [16], Alink *et al.* [17], Barbe *et al.* [13], Albrecher *et al.* [18], Biard *et al.* [10] and Kortschak and Albrecher [19], among many others.

The paper is organized as follows. In Section 2, we obtain the asymptotic finite-time ruin probabilities for regularly varying claim sizes in the risk model introduced by Boudreault *et al.* [7]. In Section 3, we derive such ruin probabilities in two other risk models, for earthquake or flooding-type risks, that take consecutive gauge-loading effects into account. The methods of proof will rely on the approximation (4) and the analysis of spacings in a conditioned Poisson process. Finally, some numerical illustrations are presented in Section 4.

2. Direct effects of each claim inter-arrival time

Our starting point is the model of Boudreault *et al.* [7] where claim amounts are of two different types depending on the length of the previous claim inter-arrival time. Hereafter, we are going to assume that the claim amounts may be dependent and they have heavy-tailed distributions.

Specifically, if an inter-arrival period V_i is larger than a fixed threshold τ , then the next claim amount X_i is given by a random variable Y_i , and if not, X_i corresponds to another random variable Z_i . As stipulated before, each vector $\mathbf{Y}^{(j)} = (Y_1, \dots, Y_j)$ is of multivariate regular variation of index $-\alpha$ and common d.f. F , while each vector $\mathbf{Z}^{(j)} = (Z_1, \dots, Z_j)$ is multivariate regularly varying of index $-\beta$ and common d.f. G .

For instance, for earthquake-type risks, one would expect that $\alpha < \beta$: the longer a period without any earthquake, the more serious will be the next earthquake. For flooding risks, the inverse situation where $\alpha > \beta$ seems to be quite plausible.

Consider the random variable $M(t, \tau)$ that gives the number of spacings of the Poisson process $\{N(s), 0 \leq s \leq t\}$ which are larger than τ . Conditioning by the number of events $N(t)$, define the following.

Let $M(n, t, \tau)$ be the random variable that counts the number, during $(0, t)$, of Poisson spacings which are larger than τ , given that $N(t) = n (\geq 1)$ and $0 < \tau < t$.

Proposition 1

If $\alpha < \beta$, for $t > 0$ and large u , the ruin probability

$$\psi(u, t) \sim \left\{ \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=1}^{\min(n, \lfloor t/\tau \rfloor)} P[M(n, t, \tau) = j] q_{j,\alpha} \right\} \bar{F}(u + ct), \tag{6}$$

while if $\alpha > \beta$, for $t > 0$ and large u ,

$$\psi(u, t) \sim \left\{ \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=\max(1, n - \lfloor t/\tau \rfloor)}^n P[M(n, t, \tau) = n - j] q_{j,\beta} \right\} \bar{G}(u + ct) \tag{7}$$

with, for $0 \leq j \leq n$,

$$P[M(n, t, \tau) = j] = \sum_{i=j}^n (-1)^{i-j} \binom{n}{j} \binom{n-j}{i-j} P\left(V_1^* > \frac{\tau}{t}, \dots, V_i^* > \frac{\tau}{t}\right), \tag{8}$$

where

$$P(V_1^* > v, \dots, V_i^* > v) = \begin{cases} 1, & v \leq 0, \\ (1 - iv)^n, & 0 < v < 1/i, \\ 0, & 1/i \leq v < 1. \end{cases} \quad (9)$$

Proof

Consider $S(t)$, the aggregate claim amount (2). Following Biard *et al.* [10], we know that if $x \mapsto P[S(t) > x]$ is regularly varying, then for large u ,

$$\psi(u, t) \sim P[S(t) > u + ct]. \quad (10)$$

To begin with, we find that

$$\begin{aligned} P[S(t) > x] &= \sum_{n=1}^{\infty} P[N(t) = n] P(X_1 + \dots + X_n > x) \\ &= \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=1}^n P[M(t, \tau) = j | N(t) = n] \\ &\quad P[X_1 + \dots + X_n > x | N(t) = n, M(t, \tau) = j] \end{aligned}$$

and by the model assumptions,

$$\begin{aligned} P[S(t) > x] &= \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=1}^n P[M(n, t, \tau) = j] \\ &\quad P(Y_1 + \dots + Y_j + Z_1 + \dots + Z_{n-j} > x). \end{aligned} \quad (11)$$

Note that

$$P[M(n, t, \tau) = j] = 0 \quad \text{for } j > t/\tau$$

so that the summation over j in (11) goes from 1 to $\min(n, \lfloor t/\tau \rfloor)$. Now, suppose that $\alpha < \beta$, so that $G(x) = o(F(x))$ for large x . Putting $S_k = Y_1 + \dots + Y_k$ and $T_k = Z_1 + \dots + Z_k$, $k \geq 1$, and using Proposition 1 in [20] and (4), we obtain

$$\begin{aligned} P(S_j + T_{n-j} > x) &\sim P(S_j > x) + P(T_{n-j} > x) \\ &\sim P(S_j > x) \sim q_{j,\alpha} \bar{F}(x), \quad 1 \leq j \leq n. \end{aligned} \quad (12)$$

Inserting (12) in (11) then yields

$$P[S(t) > x] \sim \left\{ \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=1}^{\min(n, \lfloor t/\tau \rfloor)} P[M(n, t, \tau) = j] q_{j,\alpha} \right\} \bar{F}(x);$$

that is, the function $P[S(t) > x]$ is regularly varying with parameter α . Therefore, (10) is applicable and gives the announced result (6). The approximation (7) is derived in a similar way.

Now, it remains to evaluate the p.m.f. of the random variable $M(n, t, \tau)$. As $\{N(t), t \geq 0\}$ is a Poisson process, it is well known that given $N(t) = n$, the claim instants U_1, \dots, U_n are distributed as the order statistics of n uniform random variables on $[0, t]$, and the claim inter-arrival times V_1, \dots, V_n (i.e. the associated spacings) are exchangeable random variables such that the p.d.f. of the vector (V_1^*, \dots, V_i^*) , where $V_i^* = V_i/t$, $1 \leq i \leq n$, is given by

$$\begin{aligned} f(v_1, \dots, v_i) &= \frac{n!}{(n-i)!} (1 - v_1 - \dots - v_i)^{n-i} \\ &\quad \text{for } v_1, \dots, v_i \geq 0, \quad v_1 + \dots + v_i \leq 1; \end{aligned}$$

see e.g. David and Nagaraja [21], Section 6.4. Consequently, whenever $v_1, \dots, v_i \geq 0$ with $v_1 + \dots + v_i \leq 1$,

$$P(V_1^* > v_1, \dots, V_i^* > v_i) = (1 - v_1 - \dots - v_i)^n. \quad (13)$$

In particular, choosing $v_i = \tau/t = v$ for all i in (13), we find that $P(V_1 > \tau, \dots, V_i > \tau)$ is provided by the formula (9) above. Let us consider the event that exactly j claim inter-arrival times are larger than τ , given that $N(t) = n$, with $1 \leq j \leq n$. Following e.g. David and Nagaraja [21] (p. 129–130), we then get

$$\begin{aligned} P[M(n, t, \tau) = j] &= \frac{1}{j!} \sum_{l=0}^{n-j} \frac{(-1)^l}{l!} (j+l)! \sum_{i_1 < \dots < i_{j+l}} P(V_{i_1} > \tau, \dots, V_{i_{j+l}} > \tau | N(t) = n) \\ &= \frac{1}{j!} \sum_{l=0}^{n-j} \frac{(-1)^l}{l!} (j+l)! \binom{n}{j+l} P(V_1 > \tau, \dots, V_{j+l} > \tau | N(t) = n) \\ &= \sum_{i=j}^n (-1)^{i-j} \frac{n!}{j!(i-j)!(n-i)!} P(V_1^* > v, \dots, V_i^* > v) \end{aligned} \quad (14)$$

with $v = \tau/t$. □

3. Consecutive gauge-loading effects

In this section, we want to incorporate in the risk model the observation that several consecutive claims with large, or small, inter-occurrence times are more susceptible to be followed by severe damages. By comparison with the previous model, it will be necessary this time to take (part of) the history of the claim arrival process into account. So, a gauge will be used to register the large, or small, claim inter-arrival times; initially, the gauge is empty.

3.1. Earthquake-type phenomenon

Roughly speaking, for a risk of earthquake-type, the severity of a catastrophe is expected to be more important if the latest catastrophes occur a long time ago. In this sense, let us assume that if k consecutive claim inter-arrival times are larger than τ , then a more dangerous catastrophe may arise. Moreover, just after, the gauge is put at level 0. Damages caused by standard earthquakes give multivariate regular varying vectors $\mathbf{Z}^{(j)}$ of index $-\beta$, while the more severe form regular varying vectors $\mathbf{Y}^{(j)}$ with index $-\alpha$ and common d.f. F .

Let $M_+(n, k, t, \tau)$ be the random variable that counts the number of sequences, during $(0, t)$, of k consecutive Poisson spacings which are larger than τ , given that $N(t) = n$ (≥ 1).

Proposition 2

If $\alpha < \beta$, for $t > 0$ and large u , the ruin probability

$$\psi_k^+(u, t) \sim \left\{ \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=1}^{\min(\lfloor n/k \rfloor, \lfloor t/k\tau \rfloor)} P[M_+(n, k, t, \tau) = j] q_{j, \alpha} \right\} \bar{F}(u + ct) \quad (15)$$

with, for $0 \leq j \leq \min(\lfloor n/k \rfloor, \lfloor t/k\tau \rfloor)$,

$$\begin{aligned} P[M_+(n, k, t, \tau) = j] &= \sum_{i=0}^{k-1} \sum_{x_1, \dots, x_k \in D_k} \binom{x_1 + \dots + x_k + j}{x_1, \dots, x_k, j} \\ &P[M(n, t, \tau) = n - x_1 - \dots - x_k] \bigg/ \binom{n}{x_1 + \dots + x_k}, \end{aligned} \quad (16)$$

where D_k is the set of all nonnegative integers x_1, \dots, x_k such that $\sum_{r=1}^k r x_r = n - i - kj$ and $n - \sum_{r=1}^k x_r \leq \lfloor t/\tau \rfloor$.

Proof

As in the proof of Proposition 1, we first condition by the value of $N(t)$ and then, by the value of $M_+(n, k, t, \tau)$ to get

$$\begin{aligned} P[S(t) > x] &= \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=1}^n P[M_+(n, k, t, \tau) = j] \\ &P[X_1 + \dots + X_n > x | N(t) = n, M_+(n, k, t, \tau) = j]. \end{aligned}$$

By the model assumptions and using the arguments as in the proof of Proposition 1, we have

$$P[X_1 + \dots + X_n > x | N(t) = n, M_+(n, k, t, \tau) = j] = P[Y_1 + \dots + Y_j + Z_1 + \dots + Z_{n-j} > x] \\ \sim q_{j,\alpha} \bar{F}(x), \quad 1 \leq j \leq n.$$

Thus,

$$P[S(t) > x] = \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=1}^n P[M_+(n, k, t, \tau) = j] q_{j,\alpha} \bar{F}(x),$$

which leads again to the desired approximation (15).

Now, one has to determine the p.m.f. of the random variable $M_+(n, k, t, \tau)$. This is closely related to the evaluation of the so-called binomial distributions of order k (see e.g. the book by [22]). Remember that the p.m.f. of $M(n, t, \tau)$ is given in Proposition 2.2. Denote by S the event [a Poisson spacing until time t is larger than τ , given that $N(t) = n$], and let F be the complementary event. Following [23], we observe that a typical element of the event $[M_+(n, k, t, \tau) = j]$ consists in an arrangement of the form

$$a_1 a_2 \dots a_{x_1 + \dots + x_k + j} \underbrace{SS \dots S}_i \text{ with } 0 \leq i \leq k - 1,$$

where x_1 of the a 's are F , x_2 of the a 's are SF , ..., x_k of the a 's are $\underbrace{SS \dots S}_{k-1} F$ and j of the a 's are $\underbrace{SS \dots S}_k$. Note that these x_r are subject to the constraint $x_1 + 2x_2 + \dots + kx_k + kj + i = n$. Moreover, by construction of the model, the total number of S 's has to be smaller than $\lfloor t/\tau \rfloor$, so that $x_2 + 2x_3 + (k-1)x_k + kj + i \leq \lfloor t/\tau \rfloor$. Now, such arrangements are in number

$$\binom{x_1 + \dots + x_k + j}{x_1, \dots, x_k, j}$$

and each of them has probability

$$P[M(n, t, \tau) = n - x_1 - \dots - x_k] / \binom{n}{x_1 + \dots + x_k}$$

hence the formula (16). □

3.2. Flooding-type phenomenon

For a risk of flooding-type, a close succession of claims is expected to be followed by a more important catastrophe. So, we here assume that if k consecutive claim inter-arrival times are smaller than τ , then the next claim may be more severe. Moreover, the gauge becomes empty once an inter-arrival time is larger than τ (it corresponds to a reconstruction time). Note that in the model of Section 3.1, success runs of length k are counted with a new counter just after each sequence, while in the model here, success runs of length at least k are considered with a new counter only at the first failure. In case of standard floods, damages form multivariate regular varying vectors $\mathbf{Y}^{(j)}$ of index $-\alpha$, while the more severe ones give regular varying vectors $\mathbf{Z}^{(j)}$ with index $-\beta$ and common d.f. G .

Let $M_-(n, k, t, \tau)$ be the random variable that counts the number of sequences, during $(0, t)$, of at least k consecutive Poisson spacings which are smaller than τ , given that $N(t) = n$ (≥ 1).

Proposition 3

If $\alpha > \beta$, for $t > 0$ and large u , the ruin probability

$$\psi_k^-(u, t) \sim \left\{ \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=1}^{\min(\lfloor (n+1)/(k+1) \rfloor, \lfloor t/(\tau-1) \rfloor)} P[M_-(n, k, t, \tau) = j] q_{j,\beta} \right\} \bar{G}(u + ct) \quad (17)$$

with, for $0 \leq j \leq \min(\lfloor (n+1)/(k+1) \rfloor, \lfloor t/(\tau-1) \rfloor)$,

$$P[M_-(n, k, t, \tau) = j] = \sum_{i=0}^n \sum_{x_1, \dots, x_n \in E_k} \binom{x_1 + \dots + x_n}{x_1, \dots, x_n} \\ P[M(n, t, \tau) = x_1 + \dots + x_n] / \binom{n}{x_1 + \dots + x_n}, \quad (18)$$

where E_k is the set of all nonnegative integers x_1, \dots, x_n such that $\sum_{r=1}^n r x_r = n - i$, $\sum_{r=k+1}^n x_r = j - 1_{\{i \geq k\}}$ and $\sum_{r=1}^n x_r \leq \lfloor t/\tau \rfloor$.




Figure 1. Asymptotic ruin probability $\psi(u, t)$ for the model of Section 2 as a function of the dependence parameter p in the BLL copula when $\alpha=0.5$ and $t=10$.




Figure 2. Asymptotic ruin probability $\psi(u, t)$ for the model of Section 2 as a function of the dependence parameter p in the BLL copula when $\alpha=3$ and $t=10$.

Proof

Substituting $M_-(n, k, t, \tau)$ for $M_+(n, k, t, \tau)$ and using the same way as in the proof of Proposition 2 we get, with the above assumptions, that

$$P[S(t) > x] = \sum_{n=1}^{\infty} P[N(t) = n] \sum_{j=1}^n P[M_-(n, k, t, \tau) = j] \\ P[X_1 + \dots + X_n > x | N(t) = n, M_-(n, k, t, \tau) = j]$$




Figure 3. Asymptotic ruin probability $\psi_k^+(u, t)$ for the model of Subsection 3.1 as a function of the dependence parameter p in the BLL copula when $\alpha=0.5$ and $t=20$.




Figure 4. Asymptotic ruin probability $\psi_k^+(u, t)$ for the model of Subsection 3.1 as a function of the dependence parameter p in the BLL copula when $\alpha=3$ and $t=20$.

$$\begin{aligned}
 &= \sum_{n=1}^{\infty} P[N(t)=n] \sum_{j=1}^n P[M_-(n, k, t, \tau)=j] P[Y_1 + \dots + Y_{n-j} + Z_1 + \dots + Z_j > x] \\
 &\sim \sum_{n=1}^{\infty} P[N(t)=n] \sum_{j=1}^n P[M_-(n, k, t, \tau)=j] q_{j, \beta} \bar{G}(x),
 \end{aligned}$$

which leads again to the desired approximation (17).




Figure 5. Asymptotic ruin probability $\psi(u, t)$ for the model of Section 2 as a function of τ when $\alpha=3$ and $t=10$.

To determine the p.m.f. of the random variable $M_-(n, k, t, \tau)$, one still may argue as for $M_+(n, k, t, \tau)$. Let S be here the event [a Poisson spacing until time t is smaller than τ , given that $N(t)=n$], and let F be the complementary event. An element of the event $[M_-(n, k, t, \tau)=j]$ is an arrangement of the form

$$a_1 a_2 \dots a_{x_1+\dots+x_n} \underbrace{SS\dots S}_i \text{ with } 0 \leq i \leq n,$$

where x_1 of the a 's are F , x_2 of the a 's are SF , ..., x_n of the a 's are $\underbrace{SS\dots S}_{n-1} F$, with the constraints $x_1+2x_2+\dots+nx_n+i=n$ and $x_{k+1}+\dots+x_n+1_{\{i \geq k\}}=j$. In addition, as the number of spacings larger than τ is smaller than $\lfloor t/\tau \rfloor$, one has $x_1+\dots+x_n \leq \lfloor t/\tau \rfloor$. Now, the number of such arrangements is

$$\binom{x_1+\dots+x_n}{x_1, \dots, x_n}$$

each of them having probability

$$P[M(n, t, \tau)=x_1+\dots+x_n] / \binom{n}{x_1+\dots+x_n},$$

so that the formula (18) then follows. □

4. Numerical analysis

A central step of the numerical analysis is the computation of the p.m.f. of the random variables $M(n, t, \tau)$, $M_+(n, k, t, \tau)$ and $M_-(n, k, t, \tau)$. For that, one can use the exact formulas obtained before. Another possible way is to proceed by recursion. A recursive method is provided in the Appendix for the variables M_+ and M_- (it is not simpler for M). Both methods have their own advantages and drawbacks.

- The exact formulas are easier to implement but the computation is longer.
- The recursive method works faster but implementation is fastidious.




Figure 6. Asymptotic ruin probability $\psi_k^+(u, t)$ for the model of Section 3.1 as a function of τ when $\alpha=3$ and $t=20$.

Hereafter, we have chosen to work with the exact formulas because we aim to see the impact of various dependence parameters on the asymptotic ruin probabilities. Someone who wants to investigate larger time horizons for example, might prefer to follow the recursive method.

4.1. Impact of a dependence between claim amounts

In this first part, we consider asymptotic ruin probabilities $\psi(u, t)$ for different values of $q_{k,\alpha}$ (see Table I). In order to have a spectrum of dependence, the parameter p (called ‘dependence parameter’) in the BLL copula model is allowed to vary from 0 to 1 with step 0.01. When the dependence parameter equals 0 we have the independence case, and when it equals 1 we obtain the Fréchet upper bound case. The other parameters are:

u	c	λ	Distribution of the riskier claim amount
1 000 000	10	0.1	Pareto law with parameter α

For the model of Section 2, we examine the case where $\alpha < \beta$. For $t=10$ and for both $\alpha=0.5$ and $\alpha=3$, two cases are investigated, one with $\tau=1$ and the other with $\tau=2$. Each case is plotted first separately and then in a single graph. When $\alpha=0.5$ (Figure 1) the asymptotic ruin probability is a decreasing function of the dependence parameter p . It is the opposite when $\alpha=3$ (Figure 2).

For the model of Subsection 3.1, with $k=3$, the same phenomenon is observed (see Figure 3 for $\alpha=0.5$ and Figure 4 for $\alpha=3$).

4.2. Impact of a dependence between claim inter-arrival times and claim amounts

In this second part, we examine the effect of two parameters on the asymptotic ruin probabilities: τ for the model of Section 2 and τ and k for the model of Subsection 3.1. Let us choose $\alpha=3$, for instance. For each case, we consider two dependence parameters $p=0.4$ and $p=0.8$; these are plotted separately and the relative difference is then plotted in the third graph.

For the model of Section 2 with $t=10$, as expected, the asymptotic ruin probability is a decreasing function of τ (see Figure 5). For the model of Section 3.1 with $t=20$, the asymptotic ruin probability is a decreasing function of both τ and k (see Figure 6 and Table II).

Table II. Asymptotic ruin probability $\psi_k^+(u, t)$ for the model of Section 3.1 as a function of k when $\alpha=3$ and $t=20$.				
$\alpha=3, t=20$ and $\tau=2$				
k	1	2	3	4
Ruin probability ($R1$) Dependence parameter =0.4	1.56×10^{-09}	3.08×10^{-10}	2.69×10^{-10}	6.19×10^{-11}
Ruin probability ($R2$) Dependence parameter =0.8	1.72×10^{-09}	3.21×10^{-10}	2.70×10^{-10}	6.19×10^{-11}
Relative difference $((R2 - R1)/R2)$	9.24×10^{-2}	4.01×10^{-2}	8.54×10^{-4}	1.05×10^{-6}

Appendix A

We present in this section a recursive method to compute the p.m.f. of $M_+(n, k, t, \tau)$ and $M_-(n, k, t, \tau)$. The method of proof is directly inspired from Makri and Philippou [24] (Theorem 4.1).

Proposition 4

Let

$$p_{n,n}(t) = \begin{cases} 0 & \text{if } n\tau > t, \\ \left(1 - \frac{n\tau}{t}\right)^n & \text{if } n\tau \leq t, \end{cases}$$

$$p_{0,n}(t) = 1 - p_{n,n}(t)$$

and for $1 \leq l \leq n-1$,

$$p_{l,n}(t) = \begin{cases} 0 & \text{if } l\tau > t, \\ \left(1 - \frac{l\tau}{t}\right)^n - \left(1 - \frac{(l+1)\tau}{t}\right)^n & \text{if } (l+1)\tau \leq t, \\ \left(1 - \frac{l\tau}{t}\right)^n & \text{if } l\tau \leq t < (l+1)\tau. \end{cases}$$

(a) For $n \geq k+1$ and $1 \leq j \leq \min(\lfloor n/k \rfloor, \lfloor t/(k\tau) \rfloor)$,

$$\begin{aligned} P[M_+(n, k, t, \tau) = j] &= p_{n,n}(t) \delta_{j, \lfloor n/k \rfloor} \\ &+ \frac{1}{t} \frac{p_{0,n}(t)}{B(1, n)} \int_0^\tau P[M_+(n-1, k, t-s, \tau) = j] \left(1 - \frac{s}{t}\right)^n ds \\ &+ \frac{1}{t} \sum_{l=1}^{k-1} \frac{p_{l,n}(t)}{B(l+1, n-l)} \int_{l\tau}^t \left(\frac{s}{t}\right)^l \left(1 - \frac{s}{t}\right)^{n-l-1} \\ &\times P[M_+(n-1-l, k, t-s, \tau) = j] ds \\ &+ \frac{1}{t} \sum_{l=k}^{n-1} \frac{p_{l,n}(t)}{B(l+1, n-l)} \int_{l\tau}^t \left(\frac{s}{t}\right)^l \left(1 - \frac{s}{t}\right)^{n-l-1} \\ &\times P[M_+(n-1-l, k, t-s, \tau) = j - \lfloor l/k \rfloor] ds, \end{aligned}$$

(b) for $n \geq k+1$,

$$\begin{aligned} P[M_+(n, k, t, \tau) = 0] &= \frac{1}{t} \frac{p_{0,n}(t)}{B(1, n)} \int_0^\tau P[M_+(n-1, k, t-s, \tau) = 0] \left(1 - \frac{s}{t}\right)^n ds \\ &+ \frac{1}{t} \sum_{l=1}^{k-1} \frac{p_{l,n}(t)}{B(l+1, n-l)} \int_{l\tau}^t \left(\frac{s}{t}\right)^l \left(1 - \frac{s}{t}\right)^{n-l-1} \\ &\times P[M_+(n-1-l, k, t-s, \tau) = 0] ds, \end{aligned}$$

(c) for $n \geq k + 1$ and $j > \min(\lfloor n/k \rfloor, \lfloor t/(k\tau) \rfloor)$,

$$P[M_+(n, k, t, \tau) = j] = 0,$$

(d) for $n = k$,

$$P[M_+(n, k, t, \tau) = 0] = 1 - p_{k,k}(t),$$

$$P[M_+(n, k, t, \tau) = 1] = p_{k,k}(t),$$

(e) for $0 \leq n \leq k - 1$,

$$P[M_+(n, k, t, \tau) = 0] = 1.$$

Proof

(a) Let $n \geq k + 1$ and $1 \leq j \leq \min(\lfloor n/k \rfloor, \lfloor t/(k\tau) \rfloor)$. Define, for $l = 0, \dots, n - 1$, the event $A_l = [l \text{ inter-arrival times larger than } \tau \text{ precede the first inter-arrival time smaller than } \tau \text{ in the sequence of } n \text{ inter-arrival times}]$ and the event $B = [\text{there is no inter-arrival time smaller than } \tau \text{ in the sequence of } n \text{ inter-arrival times}]$. Then, we can write that

$$[M_+(n, k, t, \tau) = j] = \left(\bigcup_{l=0}^{n-1} \{[M_+(n, k, t, \tau) = j] \cap A_l\} \right) \cup \{[M_+(n, k, t, \tau) = j] \cap B\},$$

so that we have

$$P[M_+(n, k, t, \tau) = j] = \sum_{l=0}^{n-1} P[M_+(n, k, t, \tau) = j | A_l] P(A_l) + P[M_+(n, k, t, \tau) = j | B] P(B).$$

For $l = 0$,

$$P[M_+(n, k, t, \tau) = j | A_0] = \int_0^\tau P[M_+(n-1, k, t-s, \tau) = j] f_{U_1}(s) ds.$$

For $1 \leq l \leq k - 1$,

$$P[M_+(n, k, t, \tau) = j | A_l] = \int_{l\tau}^t P[M_+(n-1-l, k, t-s, \tau) = j] f_{U_{l+1}}(s) ds.$$

For $k \leq l \leq n - 1$,

$$P[M_+(n, k, t, \tau) = j | A_l] = \int_{l\tau}^t P[M_+(n-1-l, k, t-s, \tau) = j - \lfloor l/k \rfloor] f_{U_{l+1}}(s) ds.$$

And

$$P[M_+(n, k, t, \tau) = j | B] = \delta_{j, \lfloor n/k \rfloor}.$$

From (9), we get, for $1 \leq l \leq n - 1$,

$$P(A_l) = \begin{cases} 0 & \text{if } l\tau > t, \\ P(V_1 > \tau, \dots, V_l > \tau) - P(V_1 > \tau, \dots, V_{l+1} > \tau) & \text{if } (l+1)\tau \leq t, \\ P(V_1 > \tau, \dots, V_l > \tau) & \text{if } \tau \leq t < (l+1)\tau, \end{cases}$$

$$= \begin{cases} 0 & \text{if } l\tau > t, \\ \left(1 - \frac{l\tau}{t}\right)^n - \left(1 - \frac{(l+1)\tau}{t}\right)^n & \text{if } (l+1)\tau \leq t, \\ \left(1 - \frac{l\tau}{t}\right)^n & \text{if } l\tau \leq t < (l+1)\tau, \end{cases}$$

$$P(A_0) = 1 - P(V_1 > \tau) = 1 - \left(1 - \frac{\tau}{t}\right)^n$$

and

$$P(B) = \begin{cases} 0 & \text{if } n\tau > t, \\ P(V_1 > \tau, \dots, V_n > \tau) & \text{if } n\tau \leq t, \end{cases}$$

$$= \begin{cases} 0 & \text{if } n\tau > t, \\ \left(1 - \frac{n\tau}{t}\right)^n & \text{if } n\tau \leq t. \end{cases}$$

From David and Nagaraja [21, p. 134–135], we have, for $l = 1, \dots, n$

$$f_{U_l}(u) = \frac{1}{t} \frac{1}{B(l, n+1-l)} \left(\frac{u}{t}\right)^{l-1} \left(1 - \frac{u}{t}\right)^{n-l},$$

where $B(\cdot, \cdot)$ is the Beta function. So, we deduce part (a) of the proposition.

(b) When $j = 0$ and $n \geq k$, we observe that

$$[M_+(n, k, t, \tau) = 0] = \bigcup_{l=0}^{k-1} \{[M_+(n, k, t, \tau) = 0] \cap A_l\}$$

so that

$$P[M_+(n, k, t, \tau) = 0] = \sum_{l=0}^{k-1} P[M_+(n, k, t, \tau) = 0 | A_l].$$

For $l = 0$,

$$P[M_+(n, k, t, \tau) = 0 | A_0] = \int_0^\tau P[M_+(n-1, k, t-s, \tau) = 0] f_{U_1}(s) ds.$$

For $1 \leq l \leq k-1$,

$$P[M_+(n, k, t, \tau) = 0 | A_l] = \int_{l\tau}^t P[M_+(n-1-l, k, t-s, \tau) = 0] f_{U_{l+1}}(s) ds.$$

For $l \geq k$,

$$P[M_+(n, k, t, \tau) = 0 | A_l] = 0.$$

Thus, we get part (b) of the proposition.

(c) Obvious fact.

(d) For $n = k$

$$P[M_+(n, k, t, \tau) = 1] = P(V_1 > \tau, V_2 > \tau, \dots, V_k > \tau) = \left(1 - \frac{k\tau}{t}\right)^k$$

and

$$P[M_+(n, k, t, \tau) = 0] = 1 - P[M_+(n, k, t, \tau) = 1].$$

(e) Obvious fact. □

Proposition 5

Let

$$q_{n,n}(t) = 1 - \left[n \left(1 - \frac{\tau}{t}\right)^n - \binom{n}{2} \left(1 - \frac{2\tau}{t}\right)^n + \dots + (-1)^{n-1} \left(1 - \frac{n\tau}{t}\right)^n \right],$$

$$q_{0,n}(t) = \left(1 - \frac{\tau}{t}\right)^n$$

and for $1 \leq l \leq n-1$,

$$q_{l,n}(t) = \left[(l+1) \left(1 - \frac{\tau}{t}\right)^n - \binom{l+1}{2} \left(1 - \frac{2\tau}{t}\right)^n + \dots + (-1)^l \left(1 - \frac{(l+1)\tau}{t}\right)^n \right] \\ - \left[l \left(1 - \frac{\tau}{t}\right)^n - \binom{l}{2} \left(1 - \frac{2\tau}{t}\right)^n + \dots + (-1)^{l-1} \left(1 - \frac{l\tau}{t}\right)^n \right].$$

(a) For $n \geq k+1$ and $1 \leq j \leq \min(\lfloor (n+1)/(k+1) \rfloor, \lfloor t/(\tau-1) \rfloor)$,

$$P[M_-(n, k, t, \tau) = j] = \delta_{j,1} q_{n,n}(t) \\ + \frac{1}{t} \sum_{l=0}^{k-1} \frac{q_{l,n}(t)}{B(l+1, n-l)} \int_{\tau}^t \left(\frac{s}{t}\right)^l \left(1 - \frac{s}{t}\right)^{n-l-1} \\ \times P[M_-(n-1, k, t-s, \tau) = j] ds \\ + \frac{1}{t} \sum_{l=k}^{n-1} \frac{q_{l,n}(t)}{B(l+1, n-l)} \int_{\tau}^t \left(\frac{s}{t}\right)^l \left(1 - \frac{s}{t}\right)^{n-l-1} \\ \times P[M_-(n-1-l, k, t-s, \tau) = j-1] ds.$$

(b) for $n \geq k+1$,

$$P[M_-(n, k, t, \tau) = 0] = \frac{1}{t} \sum_{l=0}^{k-1} \frac{q_{l,n}(t)}{B(l+1, n-l)} \\ \int_{t\tau}^t \left(\frac{s}{t}\right)^l \left(1 - \frac{s}{t}\right)^{n-l-1} P[M_-(n-1-l, k, t-s, \tau) = 0] ds.$$

(c) for $n \geq k+1$ and $j > \min(\lfloor (n+1)/(k+1) \rfloor, \lfloor t/(\tau-1) \rfloor)$,

$$P[M_-(n, k, t, \tau) = j] = 0.$$

(d) for $n = k$,

$$P[M_-(n, k, t, \tau) = 0] = 1 - q_{k,k}(t),$$

$$P[M_-(n, k, t, \tau) = 1] = q_{k,k}(t).$$

(e) for $0 \leq n \leq k-1$,

$$P[M_-(n, k, t, \tau) = 0] = 1.$$

Proof

We will follow an analogous argument.

(a) Let $n \geq k-1$ and $1 \leq j \leq \min(\lfloor (n+1)/(k+1) \rfloor, \lfloor t/(\tau-1) \rfloor)$. Define this time, for $l=0, \dots, n-1$, $A_l = [l \text{ inter-arrival times smaller than } \tau \text{ precede the first inter-arrival time larger than } \tau \text{ in the sequence of } n \text{ inter-arrival times}]$ and $B = [\text{there is no inter-arrival time larger than } \tau \text{ in the sequence of } n \text{ inter-arrival times}]$. Then,

$$[M_-(n, k, t, \tau) = j] = \left(\bigcup_{l=0}^{n-1} \{M_-(n, k, t, \tau) = j \cap A_l\} \right) \cup \{M_-(n, k, t, \tau) = j \cap B\}$$

so that

$$P[M_-(n, k, t, \tau) = j] = \sum_{l=0}^{n-1} P[M_-(n, k, t, \tau) = j | A_l] P(A_l) + P[M_-(n, k, t, \tau) = j | B] P(B).$$

For $0 \leq l \leq k-1$,

$$P[M_-(n, k, t, \tau) = j | A_l] = \int_{\tau}^{t-j\tau+1} P[M_-(n-1, k, t-s, \tau) = j] f_{U_{l+1}}(s) ds.$$

For $k \leq l \leq n-1$,

$$P[M_-(n, k, t, \tau) = j | A_l] = \int_{\tau}^{t-j\tau} P[M_-(n-1-l, k, t-s, \tau) = j-1] f_{U_{l+1}}(s) ds.$$

And

$$P[M_-(n, k, t, \tau) = j | B] = \delta_{j,1}.$$

From David and Nagaraja [21, 126, p. 134–135], we have, for $1 \leq l \leq n-1$,

$$\begin{aligned} P(A_l) &= P(V_1 \leq \tau, \dots, V_l \leq \tau) - P(V_1 \leq \tau, \dots, V_{l+1} \leq \tau) \\ &= P(V_1 \leq \tau \cup \dots \cup V_{l+1} \leq \tau) - P(V_1 \leq \tau \cup \dots \cup V_l \leq \tau) \\ &= \left[(l+1) \left(1 - \frac{\tau}{t}\right)^n - \binom{l+1}{2} \left(1 - \frac{2\tau}{t}\right)^n + \dots + (-1)^l \left(1 - \frac{(l+1)\tau}{t}\right)^n \right] \\ &\quad - \left[l \left(1 - \frac{\tau}{t}\right)^n - \binom{l}{2} \left(1 - \frac{2\tau}{t}\right)^n + \dots + (-1)^{l-1} \left(1 - \frac{l\tau}{t}\right)^n \right], \\ P(A_0) &= P(V_1 > \tau) = \left(1 - \frac{\tau}{t}\right)^n \end{aligned}$$

and

$$\begin{aligned} P(B) &= P(V_{(n)} \leq \tau) = 1 - P(V_{(n)} > \tau) \\ &= 1 - \left[n \left(1 - \frac{\tau}{t}\right)^n - \binom{n}{2} \left(1 - \frac{2\tau}{t}\right)^n + \dots + (-1)^{n-1} \left(1 - \frac{n\tau}{t}\right)^n \right]. \end{aligned}$$

Using again the expression of $f_{U_l}(u)$, $l = 1, \dots, n$, we then obtain part (a) of the proposition.

- (b) As for part (b) of Proposition 5.1.
- (c) Obvious fact.
- (d) For $n = k$,

$$\begin{aligned} P[M_-(n, k, t, \tau) = 0] &= P(V_{(n)} > \tau), \\ &= n \left(1 - \frac{\tau}{t}\right)^n - \binom{n}{2} \left(1 - \frac{2\tau}{t}\right)^n + \dots + (-1)^{n-1} \left(1 - \frac{n\tau}{t}\right)^n \end{aligned}$$

and

$$P[M_-(n, k, t, \tau) = 1] = 1 - P(V_{(n)} > \tau).$$

- (e) Obvious fact. □

Acknowledgements

C. Lefèvre thanks the members of the I.S.F.A for their warm welcome during his visit, and the Banque Nationale de Belgique for a partial support.

References

1. Rolski T, Schmidli H, Schmidt V, Teugels J. *Stochastic Processes for Insurance and Finance*, Wiley Series in Probability and Statistics. Wiley: Chichester, 1999.
2. Asmussen S. *Ruin Probabilities*, *Advanced Series on Statistical Science & Applied Probability*, vol. 2, World Scientific Publishing Co. Inc.: River Edge, NJ, 2000.
3. Goovaerts M, Kaas R, Dhaene J, Denuit M. *Modern Actuarial Risk Theory*. Kluwer Academic: The Netherlands, 2001.
4. Lefèvre C, Loisel S. On finite-time ruin probabilities for classical risk models. *Scandinavian Actuarial Journal* 2008; (1):41–60.
5. Albrecher H, Boxma OJ. A ruin model with dependence between claim sizes and claim intervals. *Insurance: Mathematics & Economics* 2004; 35(2):245–254.
6. Albrecher H, Teugels JL. Exponential behavior in the presence of dependence in risk theory. *Journal of Applied Probability* 2006; 43(1):257–273.

7. Boudreault M, Cossette H, Landriault D, Marceau E. On a risk model with dependence between interclaim arrivals and claim sizes. *Scandinavian Actuarial Journal* 2006; (5):265–285.
8. Meng Q, Zhang X, Guo J. On a risk model with dependence between claim sizes and claim intervals. *Statistics & Probability Letters* 2008; **78**(13):1727–1734.
9. Ambagaspitaya RS. Ultimate ruin probability in the Sparre Andersen model with dependent claim sizes and claim occurrence times. *Insurance: Mathematics and Economics* 2009; **44**(3):464–472.
10. Biard R, Lefèvre C, Loisel S. Impact of correlation crises in risk theory: asymptotics of finite-time ruin probabilities for heavy-tailed claim amounts when some independence and stationarity assumptions are relaxed. *Insurance: Mathematics and Economics* 2008; **43**(3):412–421.
11. Resnick S. The extremal dependence measure and asymptotic independence. *Stochastic Models* 2004; **20**(2):205–227.
12. Basrak B, Davis RA, Mikosch T. A characterization of multivariate regular variation. *The Annals of Applied Probability* 2002; **12**(3):908–920.
13. Barbe P, Fougères A-L, Genest C. On the tail behavior of sums of dependent risks. *Astin Bulletin* 2006; **36**(2):361–373.
14. Nelsen R. *An Introduction to Copulas*. Springer Science+Business Media, Inc: New York, 2006.
15. Wüthrich MV. Asymptotic value-at-risk estimates for sums of dependent random variables. *Astin Bulletin* 2003; **33**(1):75–92.
16. Alink S, Löwe M, Wüthrich MV. Diversification of aggregate dependent risks. *Insurance Mathematics and Economics* 2004; **35**(1):77–95.
17. Alink S, Löwe M, Wüthrich MV. Analysis of the expected shortfall of aggregate dependent risks. *Astin Bulletin* 2005; **35**(1):25–43.
18. Albrecher H, Asmussen S, Kortschak D. Tail asymptotics for the sum of two heavy-tailed dependent risks. *Extremes* 2006; **9**(2):107–130.
19. Kortschak D, Albrecher H. Asymptotic results for the sum of dependent non-identically distributed random variables. *Methodology and Computing in Applied Probability* 2009; **11**(3):279–306.
20. Cai J, Tang Q. On max-sum equivalence and convolution closure of heavy-tailed distributions and their applications. *Journal of Applied Probability* 2004; **41**(1):117–130.
21. David HA, Nagaraja HN. *Order Statistics* (3rd edn). Wiley Series in Probability and Statistics. Wiley-Interscience [John Wiley & Sons]: Hoboken, NJ, 2003.
22. Balakrishnan N, Koutras MV. *Runs and Scans with Applications*, Wiley Series in Probability and Statistics. Wiley-Interscience [John Wiley & Sons]: New York, 2002.
23. Philippou AN, Makri FS. Successes, runs and longest runs. *Statistics & Probability Letters* 1986; **4**(4):211–215.
24. Makri F, Philippou A. On binomial and circular binomial distributions of order k for l -overlapping success runs of length k . *Statistical Papers* 2005; **46**(3):411–432.