

# ANALYSIS OF A RANDOM FORESTS MODEL

G erard Biau

LSTA & LPMA<sup>1</sup>

Universit e Pierre et Marie Curie – Paris VI

Bo te 158, 175 rue du Chevaleret

75013 Paris, France

—

DMA<sup>2</sup>

Ecole Normale Sup erieure

45 rue d’Ulm

75230 Paris Cedex 05, France

gerard.biau@upmc.fr

## Abstract

Random forests are a scheme proposed by Leo Breiman in the 00’s for building a predictor ensemble with a set of decision trees that grow in randomly selected subspaces of data. Despite growing interest and practical use, there has been little exploration of the statistical properties of random forests, and little is known about the mathematical forces driving the algorithm. In this paper, we offer an in-depth analysis of a random forests model suggested by Breiman in [10], which is very close to the original algorithm. We show in particular that the procedure is consistent and adapts to sparsity, in the sense that its rate of convergence depends only on the number of strong features and not on how many noise variables are present.

*Index Terms* — Random forests, randomization, sparsity, dimension reduction, consistency, rate of convergence.

*AMS 2000 Classification:* 62G05, 62G20.

---

<sup>1</sup>Partially supported by the French “Agence Nationale pour la Recherche” under grant ANR-09-BLAN-0051-02 “CLARA”.

<sup>2</sup>Research carried out within the INRIA project “CLASSIC” hosted by Ecole Normale Sup erieure and CNRS.

# 1 Introduction

## 1.1 Random forests

In a series of papers and technical reports, Breiman [7, 8, 9, 10] demonstrated that substantial gains in classification and regression accuracy can be achieved by using ensembles of trees, where each tree in the ensemble is grown in accordance with a random parameter. Final predictions are obtained by aggregating over the ensemble. As the base constituents of the ensemble are tree-structured predictors, and since each of these trees is constructed using an injection of randomness, these procedures are called “random forests”.

Breiman’s ideas were decisively influenced by the early work of Amit and Geman [2] on geometric feature selection, the random subspace method of Ho [23] and the random split selection approach of Dietterich [19]. As highlighted by various empirical studies (see [9, 31, 18, 21] for instance), random forests have emerged as serious competitors to state of the art methods such as boosting (Freund [20]) and support vector machines (Shawe-Taylor and Cristianini [30]). They are fast and easy to implement, produce highly accurate predictions and can handle a very large number of input variables without overfitting. In fact, they are considered to be one of the most accurate general-purpose learning techniques available. The survey by Genuer et al. [21] may provide the reader with practical guidelines and a good starting point for understanding the method.

In Breiman’s approach, each tree in the collection is formed by first selecting at random, at each node, a small group of input coordinates (also called features or variables hereafter) to split on and, secondly, by calculating the best split based on these features in the training set. The tree is grown using CART methodology (Breiman et al. [11]) to maximum size, without pruning. This subspace randomization scheme is blended with bagging ([7, 13, 14, 3]) to resample, with replacement, the training data set each time a new individual tree is grown.

Although the mechanism appears simple, it involves many different driving forces which make it difficult to analyse. In fact, its mathematical properties remain to date largely unknown and, up to now, most theoretical studies have concentrated on isolated parts or stylized versions of the algorithm. Interesting attempts in this direction are by Lin and Jeon [27], who establish a connection between random forests and adaptive nearest neighbor methods (see also [4] for further results in this direction); Meinshausen [28], who studies the consistency of random forests in the context of conditional quantile prediction; and Devroye et al. [5], who offer consistency theorems for

various simplified versions of random forests and other randomized ensemble predictors. Nevertheless, the statistical mechanism of “true” random forests is not yet fully understood and is still under active investigation.

In the present paper, we go one step further into random forests by working out the properties of a model suggested by Breiman in [10]. Though this model is still simple compared to the true algorithm, it is nevertheless closer to reality than any other scheme we are aware of. The short draft [10] is essentially based on intuition and mathematical heuristics, some of them are questionable and make the document difficult to read and understand. However, the ideas presented by Breiman are worth clarifying and developing, and they will serve as a starting point for our study.

Before we formalize the model, some definitions are in order. Throughout the document, we suppose that we are given a training sample  $\mathcal{D}_n = \{(\mathbf{X}_1, Y_1), \dots, (\mathbf{X}_n, Y_n)\}$  of i.i.d.  $[0, 1]^d \times \mathbb{R}$ -valued random variables ( $d \geq 2$ ) with the same distribution as an independent generic pair  $(\mathbf{X}, Y)$  satisfying  $\mathbb{E}Y^2 < \infty$ . The space  $[0, 1]^d$  is equipped with the standard Euclidean metric. For fixed  $\mathbf{x} \in [0, 1]^d$ , our mission is to estimate the regression function  $r(\mathbf{x}) = \mathbb{E}[Y|\mathbf{X} = \mathbf{x}]$  using the data  $\mathcal{D}_n$ . In this respect, we say that a regression function estimate  $r_n(\mathbf{x})$  is consistent if  $\mathbb{E}[r_n(\mathbf{X}) - r(\mathbf{X})]^2 \rightarrow 0$  as  $n \rightarrow \infty$ . The main message of this paper is that Breiman’s procedure is consistent and adapts to sparsity, in the sense that its rate of convergence depends only on the number of strong features and not on how many noise variables are present.

## 1.2 The model

Formally, a random forest is a predictor consisting of a collection of randomized base regression trees  $\{r_n(\mathbf{x}, \Theta_m, \mathcal{D}_n), m \geq 1\}$ , where  $\Theta_1, \Theta_2, \dots$  are i.i.d. outputs of a randomizing variable  $\Theta$ . These random trees are combined to form the aggregated regression estimate

$$\bar{r}_n(\mathbf{X}, \mathcal{D}_n) = \mathbb{E}_\Theta [r_n(\mathbf{X}, \Theta, \mathcal{D}_n)],$$

where  $\mathbb{E}_\Theta$  denotes expectation with respect to the random parameter, conditionally on  $\mathbf{X}$  and the data set  $\mathcal{D}_n$ . In the following, to lighten notation a little, we will omit the dependency of the estimates in the sample, and write for example  $\bar{r}_n(\mathbf{X})$  instead of  $\bar{r}_n(\mathbf{X}, \mathcal{D}_n)$ . Note that, in practice, the above expectation is evaluated by Monte-Carlo, i.e., by generating  $M$  (usually large) random trees, and taking the average of the individual outcomes (this procedure is justified by the law of large numbers, see the appendix in Breiman [9]). The randomizing variable  $\Theta$  is used to determine how the successive

cuts are performed when building the individual trees, such as selection of the coordinate to split and position of the split.

In the model we have in mind, the variable  $\Theta$  is assumed to be independent of  $\mathbf{X}$  and the training sample  $\mathcal{D}_n$ . This excludes in particular any bootstrapping or resampling step in the training set. This also rules out any data-dependent strategy to build the trees, such as searching for optimal splits by optimizing some criterion on the actual observations. However, we allow  $\Theta$  to be based on a test sample, independent of, but distributed as,  $\mathcal{D}_n$ . This important issue will be thoroughly discussed in Section 3.

With these warnings in mind, we will assume that each individual random tree is constructed in the following way. All nodes of the tree are associated with rectangular cells such that at each step of the construction of the tree, the collection of cells associated with the leaves of the tree (i.e., external nodes) forms a partition of  $[0, 1]^d$ . The root of the tree is  $[0, 1]^d$  itself. The following procedure is then repeated  $\lceil \log_2 k_n \rceil$  times, where  $\log_2$  is the base-2 logarithm,  $\lceil \cdot \rceil$  the ceiling function and  $k_n \geq 2$  a deterministic parameter, fixed beforehand by the user, and possibly depending on  $n$ .

1. At each node, a coordinate of  $\mathbf{X} = (X^{(1)}, \dots, X^{(d)})$  is selected, with the  $j$ -th feature having a probability  $p_{nj} \in (0, 1)$  of being selected.
2. At each node, once the coordinate is selected, the split is at the midpoint of the chosen side.

Each randomized tree  $r_n(\mathbf{X}, \Theta)$  outputs the average over all  $Y_i$  for which the corresponding vector  $\mathbf{X}_i$  falls in the same cell of the random partition as  $\mathbf{X}$ . In other words, letting  $A_n(\mathbf{X}, \Theta)$  be the rectangular cell of the random partition containing  $\mathbf{X}$ ,

$$r_n(\mathbf{X}, \Theta) = \frac{\sum_{i=1}^n Y_i \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}}{\sum_{i=1}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta)},$$

where the event  $\mathcal{E}_n(\mathbf{X}, \Theta)$  is defined by

$$\mathcal{E}_n(\mathbf{X}, \Theta) = \left[ \sum_{i=1}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]} \neq 0 \right].$$

(Thus, by convention, the estimate is set to 0 on empty cells.) Taking finally expectation with respect to the parameter  $\Theta$ , the random forests regression estimate takes the form

$$\bar{r}_n(\mathbf{X}) = \mathbb{E}_{\Theta} [r_n(\mathbf{X}, \Theta)] = \mathbb{E}_{\Theta} \left[ \frac{\sum_{i=1}^n Y_i \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}}{\sum_{i=1}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta)} \right].$$

Let us now make some general remarks about this random forest model. First of all, we note that, by construction, each individual tree has exactly  $2^{\lceil \log_2 k_n \rceil}$  ( $\approx k_n$ ) terminal nodes, and each leaf has Lebesgue measure  $2^{-\lceil \log_2 k_n \rceil}$  ( $\approx 1/k_n$ ). Thus, if  $\mathbf{X}$  has uniform distribution on  $[0, 1]^d$ , there will be on average about  $n/k_n$  observations per terminal node. In particular, the choice  $k_n = n$  induces a very small number of cases in the final leaves, in accordance with the idea that the single trees should not be pruned.

Next, we see that, during the construction of the tree, at each node, each candidate coordinate  $X^{(j)}$  may be chosen with probability  $p_{nj} \in (0, 1)$ . This implies in particular  $\sum_{j=1}^d p_{nj} = 1$ . Although we do not precise for the moment the way these probabilities are generated, we stress that they may be induced by a test sample. This includes the situation where, at each node, randomness is introduced by selecting at random (with or without replacement) a small group of input features to split on, and choosing to cut the cell along the coordinate—inside this group—which most decreases some empirical criterion evaluated on the extra sample. This scheme is close to what the original random forests algorithm does, the essential difference being that the latter algorithm uses the actual data set to calculate the best splits. This point will be properly discussed in Section 3.

Finally, the requirement that the splits are always achieved at the middle of the cell sides is mainly technical, and it could eventually be replaced by a more involved random mechanism—based on the test sample, at the price of a much more complicated analysis.

The document is organized as follows. In Section 2, we prove that the random forests regression estimate  $\bar{r}_n$  is consistent and discuss its rate of convergence. As a striking result, we show under a sparsity framework that the rate of convergence depends only on the number of active (or strong) variables and not on the dimension of the ambient space. This feature is particularly desirable in high-dimensional regression, when the number of variables can be much larger than the sample size, and may explain why random forests are able to handle a very large number of input variables without overfitting. Section 3 is devoted to a discussion, and proofs are postponed to Section 4.

## 2 Consistency results

Throughout the document, we will denote by  $N_n(\mathbf{X}, \Theta)$  the number of data points falling in the same cell as  $\mathbf{X}$ , i.e.,

$$N_n(\mathbf{X}, \Theta) = \sum_{i=1}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}.$$

We start the analysis with the following simple theorem, which shows that the random forests estimate  $\bar{r}_n$  is consistent.

**Theorem 2.1** *Assume that the distribution of  $\mathbf{X}$  has support on  $[0, 1]^d$ . Then the random forests estimate  $\bar{r}_n$  is consistent whenever  $p_{nj} \log k_n \rightarrow \infty$  for all  $j = 1, \dots, d$  and  $k_n/n \rightarrow 0$  as  $n \rightarrow \infty$ .*

Theorem 2.1 mainly serves as an illustration of how the consistency problem of random forests predictors may be attacked. It encompasses, in particular, the situation where, at each node, the coordinate to split is chosen uniformly at random over the  $d$  candidates. In this “purely random” model,  $p_{nj} = 1/d$ , independently of  $n$  and  $j$ , and consistency is ensured as long as  $k_n \rightarrow \infty$  and  $k_n/n \rightarrow 0$ . This is however a radically simplified version of the random forests used in practice, which does not explain the good performance of the algorithm. To achieve this goal, a more in-depth analysis is needed.

There is empirical evidence that many signals in high-dimensional spaces admit a sparse representation. As an example, wavelet coefficients of images often exhibit exponential decay, and a relatively small subset of all wavelet coefficients allow for a good approximation of the original image. Such signals have few non-zero coefficients and can therefore be described as sparse in the signal domain (see for instance [12]). Similarly, recent advances in high-throughput technologies such as microarrays and aCGH arrays indicate that, despite the huge dimensionality of problems, only a small number of genes may play a role in determining the outcome and be required to create good predictors ([33] for instance). Sparse estimation is playing an increasingly important role in the statistics and machine learning communities, and several methods have recently been developed in both fields, which rely upon the notion of sparsity (e.g. penalty methods like the Lasso and Dantzig selector, see [32, 16, 15, 6] and the references therein).

Following this idea, we will assume in our setting that the target regression function  $r(\mathbf{X}) = \mathbb{E}[Y|\mathbf{X}]$ , which is initially a function of  $\mathbf{X} = (X^{(1)}, \dots, X^{(d)})$ , depends in fact only on a nonempty subset  $\mathcal{S}$  (for **S**trong) of the  $d$  features. In other words, letting  $\mathbf{X}_{\mathcal{S}} = (X_j : j \in \mathcal{S})$  and  $S = \text{Card } \mathcal{S}$ , we have

$$r(\mathbf{X}) = \mathbb{E}[Y|\mathbf{X}_{\mathcal{S}}]$$

or equivalently, for any  $\mathbf{x} \in [0, 1]^d$ ,

$$r(\mathbf{x}) = r^*(\mathbf{x}_{\mathcal{S}}) \quad \mu\text{-a.s.}, \quad (2.1)$$

where  $\mu$  is the distribution of  $\mathbf{X}$  and  $r^* : [0, 1]^S \rightarrow \mathbb{R}$  is the section of  $r$  corresponding to  $\mathcal{S}$ . To avoid trivialities, we will assume throughout that  $\mathcal{S}$

is nonempty, with  $S \geq 2$ . The variables in the set  $\mathcal{W} = \{1, \dots, d\} - \mathcal{S}$  (for *Weak*) have thus no influence on the response and could be safely removed. In the dimension reduction scenario we have in mind, the ambient dimension  $d$  can be very large, much larger than the sample size  $n$ , but we believe that the representation is sparse, i.e., that very few coordinates of  $r$  are non-zero, with indices corresponding to the set  $\mathcal{S}$ . Note however that representation (2.1) does not forbid the somehow undesirable case where  $S = d$ . As such, the value  $S$  characterizes the sparsity of the model: the smaller  $S$ , the sparser  $r$ .

Within this sparsity framework, it is intuitively clear that the coordinate-sampling probabilities should ideally satisfy the constraints  $p_{nj} = 1/S$  for  $j \in \mathcal{S}$  (and, consequently,  $p_{nj} = 0$  otherwise). However, this is a too strong requirement, which has no chance to be satisfied in practice, except maybe in some special situations where we know beforehand which variables are important and which are not. Thus, to stick to reality, we will rather require in the following that  $p_{nj} = (1/S)(1 + \xi_{nj})$  for  $j \in \mathcal{S}$  (and  $p_{nj} = \xi_{nj}$  otherwise), where  $p_{nj} \in (0, 1)$  and each  $\xi_{nj}$  tends to 0 as  $n$  tends to infinity. We will see in Section 3 how to design a randomization mechanism to obtain such probabilities, on the basis of a test sample independent of the training set  $\mathcal{D}_n$ .

We have now enough material for a deeper understanding of the random forests algorithm. To lighten notation a little, we will write

$$W_{ni}(\mathbf{X}, \Theta) = \frac{\mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}}{N_n(\mathbf{X}, \Theta)} \mathbf{1}_{\mathcal{E}_n(\mathbf{x}, \Theta)},$$

so that the estimate takes the form

$$\bar{r}_n(\mathbf{X}) = \sum_{i=1}^n \mathbb{E}_{\Theta} [W_{ni}(\mathbf{X}, \Theta)] Y_i.$$

Let us start with the variance/bias decomposition

$$\mathbb{E} [\bar{r}_n(\mathbf{X}) - r(\mathbf{X})]^2 = \mathbb{E} [\bar{r}_n(\mathbf{X}) - \tilde{r}_n(\mathbf{X})]^2 + \mathbb{E} [\tilde{r}_n(\mathbf{X}) - r(\mathbf{X})]^2, \quad (2.2)$$

where we set

$$\tilde{r}_n(\mathbf{X}) = \sum_{i=1}^n \mathbb{E}_{\Theta} [W_{ni}(\mathbf{X}, \Theta)] r(\mathbf{X}_i).$$

The two terms of (2.2) will be examined separately, in Proposition 2.1 and Proposition 2.2, respectively. Throughout, the symbol  $\mathbb{V}$  denotes variance.

**Proposition 2.1** Assume that  $\mathbf{X}$  is uniformly distributed on  $[0, 1]^d$  and, for all  $\mathbf{x} \in \mathbb{R}^d$ ,

$$\sigma^2(\mathbf{x}) = \mathbb{V}[Y | \mathbf{X} = \mathbf{x}] \leq \sigma^2$$

for some positive constant  $\sigma^2$ . Then, if  $p_{nj} = (1/S)(1 + \xi_{nj})$  for  $j \in \mathcal{S}$ ,

$$\mathbb{E} [\bar{r}_n(\mathbf{X}) - \tilde{r}_n(\mathbf{X})]^2 \leq C\sigma^2 \left( \frac{S^2}{S-1} \right)^{S/2d} (1 + \xi_n) \frac{k_n}{n(\log k_n)^{S/2d}},$$

where

$$C = \frac{288}{\pi} \left( \frac{\pi \log 2}{16} \right)^{S/2d}.$$

The sequence  $(\xi_n)$  depends on the sequences  $\{(\xi_{nj}) : j \in \mathcal{S}\}$  only and tends to 0 as  $n$  tends to infinity.

**Remark 1** A close inspection of the end of the proof of Proposition 2.1 reveals that

$$1 + \xi_n = \prod_{j \in \mathcal{S}} \left[ (1 + \xi_{nj})^{-1} \left( 1 - \frac{\xi_{nj}}{S-1} \right)^{-1} \right]^{1/2d}.$$

In particular, if  $a < p_{nj} < b$  for some constants  $a, b \in (0, 1)$ , then

$$1 + \xi_n \leq \left( \frac{S-1}{S^2 a(1-b)} \right)^{S/2d}.$$

■

The main message of Proposition 2.1 is that the variance of the forests estimate is  $\mathcal{O}(k_n/(n(\log k_n)^{S/2d}))$ . This result is interesting by itself since it shows the effect of aggregation on the variance of the forest. To understand this remark, recall that individual (random or not) trees are proved to be consistent by letting the number of cases in each terminal node become large (see [17, Chapter 20]), with a typical variance of the order  $k_n/n$ . Thus, for such trees, the choice  $k_n = n$  (i.e., about one observation on average in each terminal node) is clearly not suitable and leads to serious overfitting and variance explosion. On the other hand, the variance of the forest is of the order  $k_n/(n(\log k_n)^{S/2d})$ . Therefore, letting  $k_n = n$ , the variance is of the order  $1/(\log n)^{S/2d}$ , a quantity which still goes to 0 as  $n$  grows! Proof of Proposition 2.1 reveals that this log term is a by-product of the  $\Theta$ -averaging process, which appears by taking into consideration the correlation between trees. We believe that it provides an interesting perspective on why random

forests are still able to do a good job, despite the fact that individual trees are not pruned.

Note finally that the requirement that  $\mathbf{X}$  is uniformly distributed on the hypercube could be safely replaced by the assumption that  $\mathbf{X}$  has a density with respect to the Lebesgue measure on  $[0, 1]^d$  and the density is bounded above and below. The case where the density of  $\mathbf{X}$  is not bounded from below necessitates a specific analysis, which we believe is beyond the scope of the present paper. We refer the reader to [4] for results in this direction.

Let us now turn to the analysis of the bias term in equality (2.2).

**Proposition 2.2** *Assume that  $\mathbf{X}$  is uniformly distributed on  $[0, 1]^d$  and  $r$  is  $L$ -Lipschitz on  $[0, 1]^d$ . Then, if  $p_{nj} = (1/S)(1 + \xi_{nj})$  for  $j \in \mathcal{S}$ ,*

$$\mathbb{E} [\tilde{r}_n(\mathbf{X}) - r(\mathbf{X})]^2 \leq \frac{2SL^2}{k_n^{\frac{0.75}{S \log 2}(1+\gamma_n)}} + \left[ \sup_{\mathbf{x} \in [0,1]^d} r^2(\mathbf{x}) \right] e^{-n/2k_n},$$

where  $\gamma_n = \min_{j \in \mathcal{S}} \xi_{nj}$  tends to 0 as  $n$  tends to infinity.

This result essentially shows that the rate at which the bias decreases to 0 depends on the number of strong variables, not on  $d$ . In particular, the quantity  $k_n^{-(0.75/(S \log 2))(1+\gamma_n)}$  should be compared with the ordinary partitioning estimate bias, which is of the order  $k_n^{-2/d}$  under the smoothness conditions of Proposition 2.2 (see for instance [22]). In this respect, it is easy to see that  $k_n^{-(0.75/(S \log 2))(1+\gamma_n)} = o(k_n^{-2/d})$  as soon as  $S \leq \lfloor 0.54d \rfloor$  ( $\lfloor \cdot \rfloor$  is the integer part function). In other words, when the number of active variables is less than (roughly) half of the ambient dimension, the bias of the random forests regression estimate decreases to 0 much faster than the usual rate. The restriction  $S \leq \lfloor 0.54d \rfloor$  is not severe, since in all practical situations we have in mind,  $d$  is usually very large with respect to  $S$  (this is, for instance, typically the case in modern genome biology problems, where  $d$  may be of the order of billions, and in any case much larger than the actual number of active features). Note at last that, contrary to Proposition 2.1, the term  $e^{-n/2k_n}$  prevents the extreme choice  $k_n = n$  (one observation on average in each terminal node). Indeed, an inspection of the proof of Proposition 2.2 reveals that this exponentially small term accounts for the probability that  $N_n(\mathbf{X}, \Theta)$  is precisely 0, i.e.,  $A_n(\mathbf{X}, \Theta)$  is empty.

Recalling the elementary inequality  $ze^{-nz} \leq e^{-1}/n$  for  $z \in [0, 1]$ , we may finally join Proposition 2.1 and Proposition 2.2 and state our main theorem.

**Theorem 2.2** Assume that  $\mathbf{X}$  is uniformly distributed on  $[0, 1]^d$ ,  $r$  is  $L$ -Lipschitz on  $[0, 1]^d$  and, for all  $\mathbf{x} \in \mathbb{R}^d$ ,

$$\sigma^2(\mathbf{x}) = \mathbb{V}[Y | \mathbf{X} = \mathbf{x}] \leq \sigma^2$$

for some positive constant  $\sigma^2$ . Then, if  $p_{nj} = (1/S)(1 + \xi_{nj})$  for  $j \in \mathcal{S}$ , letting  $\gamma_n = \min_{j \in \mathcal{S}} \xi_{nj}$ , we have

$$\mathbb{E} [\bar{r}_n(\mathbf{X}) - r(\mathbf{X})]^2 \leq \Xi_n \frac{k_n}{n} + \frac{2SL^2}{k_n^{\frac{0.75}{S \log 2}(1+\gamma_n)}},$$

where

$$\Xi_n = C\sigma^2 \left( \frac{S^2}{S-1} \right)^{S/2d} (1 + \xi_n) + 2e^{-1} \left[ \sup_{\mathbf{x} \in [0,1]^d} r^2(\mathbf{x}) \right]$$

and

$$C = \frac{288}{\pi} \left( \frac{\pi \log 2}{16} \right)^{S/2d}.$$

The sequence  $(\xi_n)$  depends on the sequences  $\{(\xi_{nj}) : j \in \mathcal{S}\}$  only and tends to 0 as  $n$  tends to infinity.

As we will see in Section 3, it may be safely assumed that the randomization process allows for  $\xi_{nj} \log n \rightarrow 0$  as  $n \rightarrow \infty$ , for all  $j \in \mathcal{S}$ . Thus, under this condition, Theorem 2.2 shows that with the optimal choice

$$k_n \propto n^{1/(1+\frac{0.75}{S \log 2})},$$

we get

$$\mathbb{E} [\bar{r}_n(\mathbf{X}) - r(\mathbf{X})]^2 = \mathcal{O} \left( n^{\frac{-0.75}{S \log 2 + 0.75}} \right).$$

This result can be made more precise. Denote by  $\mathcal{F}$  the class of  $(L, \sigma^2)$ -smooth distributions  $(\mathbf{X}, Y)$  such that  $\mathbf{X}$  has uniform distribution on  $[0, 1]^d$ , the regression function  $r$  is Lipschitz with constant  $L$  and, for all  $\mathbf{x} \in \mathbb{R}^d$ ,  $\sigma^2(\mathbf{x}) = \mathbb{V}[Y | \mathbf{X} = \mathbf{x}] \leq \sigma^2$ .

**Corollary 2.1** Let

$$\Xi = C\sigma^2 \left( \frac{S^2}{S-1} \right)^{S/2d} + 2e^{-1} \left[ \sup_{\mathbf{x} \in [0,1]^d} r^2(\mathbf{x}) \right]$$

and

$$C = \frac{288}{\pi} \left( \frac{\pi \log 2}{16} \right)^{S/2d}.$$

Then, if  $p_{nj} = (1/S)(1 + \xi_{nj})$  for  $j \in \mathcal{S}$ , with  $\xi_{nj} \log n \rightarrow 0$  as  $n \rightarrow \infty$ , for the choice

$$k_n \propto \left( \frac{L^2}{\Xi} \right)^{1/(1 + \frac{0.75}{S \log 2})} n^{1/(1 + \frac{0.75}{S \log 2})},$$

we have

$$\limsup_{n \rightarrow \infty} \sup_{(\mathbf{X}, Y) \in \mathcal{F}} \frac{\mathbb{E} [\bar{r}_n(\mathbf{X}) - r(\mathbf{X})]^2}{\left( \Xi L^{\frac{2S \ln 2}{0.75}} \right)^{\frac{0.75}{S \ln 2 + 0.75}} n^{\frac{-0.75}{S \log 2 + 0.75}}} \leq \Lambda,$$

where  $\Lambda$  is a positive constant independent of  $r$ ,  $L$  and  $\sigma^2$ .

This result reveals the fact that the  $L_2$ -rate of convergence of  $\bar{r}_n(\mathbf{X})$  to  $r(\mathbf{X})$  depends only on the number  $S$  of strong variables, and not on the ambient dimension  $d$ . The main message of Corollary 2.1 is that if we are able to properly tune the probability sequences  $(p_{nj})_{n \geq 1}$  and make them sufficiently fast to track the informative features, then the rate of convergence of the random forests estimate will be of the order  $n^{\frac{-0.75}{S \log 2 + 0.75}}$ . This rate is strictly faster than the usual rate  $n^{-2/(d+2)}$  as soon as  $S \leq \lfloor 0.54d \rfloor$ . To understand this point, just recall that the rate  $n^{-2/(d+2)}$  is minimax optimal for the considered smoothness class  $\mathcal{F}$  (see for example Ibragimov and Khasminskii [24, 25, 26]), seen as a collection of regression functions over  $[0, 1]^d$ . However, in our setting, the intrinsic dimension of the regression problem is  $S$ , not  $d$ , and the random forests estimate cleverly adapts to the sparsity of the problem. As an illustration, Figure 1 shows the plot of the function  $S \mapsto 0.75/(S \log 2 + 0.75)$  for  $S$  ranging from 2 to  $d = 100$ .

**Remark 2** The reduced-dimensional rate  $n^{\frac{-0.75}{S \log 2 + 0.75}}$  is strictly larger than the  $S$ -dimensional optimal rate  $n^{-2/(S+2)}$ . We do not know whether the latter rate can be achieved by the algorithm. ■

**Remark 3** The optimal parameter  $k_n$  of Corollary 2.1 depends on the unknown distribution of  $(\mathbf{X}, Y)$ , especially on the smoothness of the regression function and the effective dimension  $S$ . To correct this situation, adaptive (i.e., data-dependent) choices of  $k_n$ , such as data-splitting or cross-validation, should preserve the rate of convergence of the estimate. Another route we may follow is to analyse the effect of bootstrapping the sample before growing the individual trees (i.e., bagging). It is our belief that this procedure should also preserve the rate of convergence, even for overfitted trees ( $k_n \approx n$ ), in the spirit of [3]. However, such a study is beyond the scope of the present paper. ■

**Remark 4** For further references, it is interesting to note that Proposition 2.1 (variance term) is a consequence of aggregation, whereas Proposition 2.2 (bias term) is a consequence of randomization. ■




Figure 1: Plot of the function  $S \mapsto 0.75/(S \log 2 + 0.75)$  for  $S$  ranging from 2 to  $d = 100$ . The horizontal line shows the value of the  $d$ -dimensional rate power  $2/(d + 2) \approx 0.0196$ .

### 3 Discussion

The results which have been obtained in Section 2 rely on appropriate behavior of the probability sequences  $(p_{nj})_{n \geq 1}$ ,  $j = 1, \dots, d$ . We recall that these sequences should be in  $(0, 1)$  and obey the constraints  $p_{nj} = (1/S)(1 + \xi_{nj})$  for  $j \in \mathcal{S}$  (and  $p_{nj} = \xi_{nj}$  otherwise), where the  $(\xi_{nj})_{n \geq 1}$  tend to 0 as  $n$  tends to infinity. In other words, at each step of the construction of the individual trees, the random procedure should track and preferentially cut the strong coordinates. In this more informal section, we briefly discuss a random mechanism for inducing such probability sequences.

Suppose, to start with an imaginary scenario, that we already know which coordinates are strong, and which are not. In this ideal case, the random selection procedure described in the introduction may be easily made more precise as follows. A positive integer  $M_n$ —possibly depending on  $n$ —is fixed beforehand and the following splitting scheme is iteratively repeated at each node of the tree:

1. Select at random, with replacement,  $M_n$  candidate coordinates to split on.

2. If the selection is all weak, then choose one at random to split on. If there is more than one strong variable elected, choose one at random and cut.

Within this framework, it is easy to see that each coordinate in  $\mathcal{S}$  will be cut with the “ideal” probability

$$p_n^* = \frac{1}{S} \left[ 1 - \left( 1 - \frac{S}{d} \right)^{M_n} \right].$$

Though this is an idealized model, it already gives some information about the choice of the parameter  $M_n$ , which, in accordance with the results of Section 2 (Corollary 2.1), should satisfy

$$\left( 1 - \frac{S}{d} \right)^{M_n} \log n \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This is true as soon as

$$M_n \rightarrow \infty \quad \text{and} \quad \frac{M_n}{\log n} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

This result is consistent with the general empirical finding that  $M_n$  does not need to be very large (see, for example, Breiman [9]), but not with the widespread belief that  $M_n$  should not depend on  $n$ . Note also that if the  $M_n$  features are chosen at random *without* replacement, then things are even more simple since, in this case,  $p_n^* = 1/S$  for all  $n$  large enough.

In practice, we have only a vague idea about the size and content of the set  $\mathcal{S}$ . However, to circumvent this problem, we may use the observations of an independent test set  $\mathcal{D}'_n$  (say, of the same size as  $\mathcal{D}_n$ ) in order to mimic the ideal split probability  $p_n^*$ . To illustrate this mechanism, suppose—to keep things simple—that the model is linear, i.e.,

$$Y = \sum_{j \in \mathcal{S}} a_j X^{(j)} + \varepsilon,$$

where  $\mathbf{X} = (X^{(1)}, \dots, X^{(d)})$  is uniformly distributed over  $[0, 1]^d$ , the  $a_j$  are non-zero real numbers, and  $\varepsilon$  is a zero-mean random noise, which is assumed to be independent of  $\mathbf{X}$  and with finite variance. Note that, in accordance with our sparsity assumption,  $r(\mathbf{X}) = \sum_{j \in \mathcal{S}} a_j X^{(j)}$  depends on  $\mathbf{X}_{\mathcal{S}}$  only.

Assume now that we have done some splitting and arrived at a current set of terminal nodes. Consider any of these nodes, say  $A = \prod_{j=1}^d A_j$ , fix a

coordinate  $j \in \{1, \dots, d\}$ , and look at the weighted conditional variance  $\mathbb{V}[Y|X^{(j)}] \mathbb{P}(X^{(j)} \in A_j)$ . It is a simple exercise to prove that if  $\mathbf{X}$  is uniform and  $j \in \mathcal{S}$ , then the split on the  $j$ -th side which most decreases the weighted conditional variance is at the midpoint of the node, with a variance decrease equal to  $a_j^2/16 > 0$ . On the other hand, if  $j \in \mathcal{W}$ , the decrease of the variance is always 0, whatever the location of the split.

On the practical side, the conditional variances are of course unknown, but they may be estimated by replacing the theoretical quantities by their respective sample estimates (as in the CART procedure, see Breiman et al. [9, Chapter 8] for a thorough discussion) evaluated on the test sample  $\mathcal{D}'_n$ . This suggests the following procedure, at each node of the tree:

1. Select at random, with replacement,  $M_n$  candidate coordinates to split on.
2. For each of the  $M_n$  elected coordinates, calculate the best split, i.e., the split which most decreases the within-node sum of squares on the test sample  $\mathcal{D}'_n$ .
3. Select one variable at random among the coordinates which output the best within-node sum of squares decreases, and cut.

This procedure is indeed close to what the random forests algorithm does. The essential difference is that we suppose to have at hand a second test sample  $\mathcal{D}'_n$ , whereas the original algorithm performs the search of the optimal cuts on the original observations  $\mathcal{D}_n$ . This point is important, since the use of an extra sample preserves the independence of  $\Theta$  (the random mechanism) and  $\mathcal{D}_n$  (the training sample). We do not know whether our results are still true if  $\Theta$  depends on  $\mathcal{D}_n$ , but the analysis does not appear to be simple. Note also that, at step 3, a threshold (or a test procedure, as suggested in Amaratunga et al. [1]) should be used to choose among the most significant variables, whereas the actual algorithm just selects the best one.

This empirical randomization scheme leads to complicate probabilities of cuts which, this time, vary at each node of each tree and are not easily amenable to analysis. Nevertheless, observing that the average number of cases per terminal node is about  $n/k_n$ , it may be inferred by the law of large numbers that each variable in  $\mathcal{S}$  will be cut with probability

$$p_{nj} \approx \frac{1}{S} \left[ 1 - \left( 1 - \frac{S}{d} \right)^{M_n} \right] (1 + \zeta_{nj}),$$

where  $\zeta_{nj}$  is of the order  $\mathcal{O}(k_n/n)$ , a quantity which anyway goes fast to 0 as  $n$  tends to infinity. Put differently, for  $j \in \mathcal{S}$ ,

$$p_{nj} \approx \frac{1}{S} (1 + \xi_{nj}),$$

where  $\xi_{nj}$  goes to 0 and satisfies the constraint  $\xi_{nj} \log n \rightarrow 0$  as  $n$  tends to infinity, provided  $k_n \log n/n \rightarrow 0$ ,  $M_n \rightarrow \infty$  and  $M_n/\log n \rightarrow \infty$ . This is coherent with the requirements of Corollary 2.1. We realize however that this is a rough approach, and that more theoretical work is needed here to fully understand the mechanisms involved in Breiman's original randomization process.

## 4 Proofs

Throughout this section, we will make repeated use of the following two facts.

**Fact 4.1** *Let  $K_{nj}(\mathbf{X}, \Theta)$  be the number of times the terminal node  $A_n(\mathbf{X}, \Theta)$  is split on the  $j$ -th coordinate ( $j = 1, \dots, d$ ). Then, conditionally on  $\mathbf{X}$ ,  $K_{nj}(\mathbf{X}, \Theta)$  has binomial distribution with parameters  $\lceil \log_2 k_n \rceil$  and  $p_{nj}$  (by independence of  $\mathbf{X}$  and  $\Theta$ ). Moreover, by construction,*

$$\sum_{j=1}^d K_{nj}(\mathbf{X}, \Theta) = \lceil \log_2 k_n \rceil.$$

Recall that we denote by  $N_n(\mathbf{X}, \Theta)$  the number of data points falling in the same cell as  $\mathbf{X}$ , i.e.,

$$N_n(\mathbf{X}, \Theta) = \sum_{i=1}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}.$$

Let  $\lambda$  be the Lebesgue measure on  $[0, 1]^d$ .

**Fact 4.2** *By construction,*

$$\lambda(A_n(\mathbf{X}, \Theta)) = 2^{-\lceil \log_2 k_n \rceil}.$$

*In particular, if  $\mathbf{X}$  is uniformly distributed on  $[0, 1]^d$ , then the distribution of  $N_n(\mathbf{X}, \Theta)$  conditionally on  $\mathbf{X}$  and  $\Theta$  is binomial with parameters  $n$  and  $2^{-\lceil \log_2 k_n \rceil}$  (by independence of the random variables  $\mathbf{X}, \mathbf{X}_1, \dots, \mathbf{X}_n, \Theta$ ).*

## 4.1 Proof of Theorem 2.1

Observe first that, by Jensen's inequality,

$$\begin{aligned}\mathbb{E} [\bar{r}_n(\mathbf{X}) - r(\mathbf{X})]^2 &= \mathbb{E} [\mathbb{E}_\Theta [r_n(\mathbf{X}, \Theta) - r(\mathbf{X})]]^2 \\ &\leq \mathbb{E} [r_n(\mathbf{X}, \Theta) - r(\mathbf{X})]^2.\end{aligned}$$

A slight adaptation of Theorem 4.2 in Györfi et al. [22] shows that  $\bar{r}_n$  is consistent if both  $\text{diam}(A_n(\mathbf{X}, \Theta)) \rightarrow 0$  in probability and  $N_n(\mathbf{X}, \Theta) \rightarrow \infty$  in probability.

Let us first prove that  $N_n(\mathbf{X}, \Theta) \rightarrow \infty$  in probability. To see this, consider the random tree partition defined by  $\Theta$ , which has by construction exactly  $2^{\lceil \log_2 k_n \rceil}$  rectangular cells, say  $A_1, \dots, A_{2^{\lceil \log_2 k_n \rceil}}$ . Let  $N_1, \dots, N_{2^{\lceil \log_2 k_n \rceil}}$  denote the number of observations among  $\mathbf{X}, \mathbf{X}_1, \dots, \mathbf{X}_n$  falling in these  $2^{\lceil \log_2 k_n \rceil}$  cells, and let  $\mathcal{C} = \{\mathbf{X}, \mathbf{X}_1, \dots, \mathbf{X}_n\}$  denote the set of positions of these  $n + 1$  points. Since these points are independent and identically distributed, fixing the set  $\mathcal{C}$  and  $\Theta$ , the conditional probability that  $\mathbf{X}$  falls in the  $\ell$ -th cell equals  $N_\ell / (n + 1)$ . Thus, for every fixed  $M \geq 0$ ,

$$\begin{aligned}\mathbb{P}(N_n(\mathbf{X}, \Theta) < M) &= \mathbb{E} [\mathbb{P}(N_n(\mathbf{X}, \Theta) < M \mid \mathcal{C}, \Theta)] \\ &= \mathbb{E} \left[ \sum_{\ell=1, \dots, 2^{\lceil \log_2 k_n \rceil}: N_\ell < M} \frac{N_\ell}{n + 1} \right] \\ &\leq \frac{M 2^{\lceil \log_2 k_n \rceil}}{n + 1} \\ &\leq \frac{2Mk_n}{n + 1},\end{aligned}$$

which converges to 0 by our assumption on  $k_n$ .

It remains to show that  $\text{diam}(A_n(\mathbf{X}, \Theta)) \rightarrow 0$  in probability. To this aim, let  $V_{nj}(\mathbf{X}, \Theta)$  be the size of the  $j$ -th dimension of the rectangle containing  $\mathbf{X}$ . Clearly, it suffices to show that  $V_{nj}(\mathbf{X}, \Theta) \rightarrow 0$  in probability for all  $j = 1, \dots, d$ . To this end, note that

$$V_{nj}(\mathbf{X}, \Theta) \stackrel{D}{=} 2^{-K_{nj}(\mathbf{X}, \Theta)},$$

where, conditionally on  $\mathbf{X}$ ,  $K_{nj}(\mathbf{X}, \Theta)$  has a binomial  $\mathcal{B}(\lceil \log_2 k_n \rceil, p_{nj})$  distribution, representing the number of times the box containing  $\mathbf{X}$  is split along the  $j$ -th coordinate (Fact 4.1). Thus

$$\begin{aligned}\mathbb{E} [V_{nj}(\mathbf{X}, \Theta)] &= \mathbb{E} [2^{-K_{nj}(\mathbf{X}, \Theta)}] \\ &= \mathbb{E} [\mathbb{E} [2^{-K_{nj}(\mathbf{X}, \Theta)} \mid \mathbf{X}]] \\ &= (1 - p_{nj}/2)^{\lceil \log_2 k_n \rceil},\end{aligned}$$

which tends to 0 as  $p_{nj} \log k_n \rightarrow \infty$ .

## 4.2 Proof of Proposition 2.1

Recall that

$$\bar{r}_n(\mathbf{X}) = \sum_{i=1}^n \mathbb{E}_{\Theta} [W_{ni}(\mathbf{X}, \Theta)] Y_i,$$

where

$$W_{ni}(\mathbf{X}, \Theta) = \frac{\mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}}{N_n(\mathbf{X}, \Theta)} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta)}$$

and

$$\mathcal{E}_n = [N_n(\mathbf{X}, \Theta) \neq 0].$$

Similarly,

$$\tilde{r}_n(\mathbf{X}) = \sum_{i=1}^n \mathbb{E}_{\Theta} [W_{ni}(\mathbf{X}, \Theta)] r(\mathbf{X}_i).$$

We have

$$\begin{aligned} \mathbb{E} [\bar{r}_n(\mathbf{X}) - \tilde{r}_n(\mathbf{X})]^2 &= \mathbb{E} \left[ \sum_{i=1}^n \mathbb{E}_{\Theta} [W_{ni}(\mathbf{X}, \Theta)] (Y_i - r(\mathbf{X}_i)) \right]^2 \\ &= \mathbb{E} \left[ \sum_{i=1}^n \mathbb{E}_{\Theta}^2 [W_{ni}(\mathbf{X}, \Theta)] (Y_i - r(\mathbf{X}_i))^2 \right] \\ &\quad \text{(the cross terms are 0 since } \mathbb{E}[Y_i | \mathbf{X}_i] = r(\mathbf{X}_i)\text{)} \\ &= \mathbb{E} \left[ \sum_{i=1}^n \mathbb{E}_{\Theta}^2 [W_{ni}(\mathbf{X}, \Theta)] \sigma^2(\mathbf{X}_i) \right] \\ &\leq \sigma^2 \mathbb{E} \left[ \sum_{i=1}^n \mathbb{E}_{\Theta}^2 [W_{ni}(\mathbf{X}, \Theta)] \right] \\ &= n\sigma^2 \mathbb{E} [\mathbb{E}_{\Theta}^2 [W_{n1}(\mathbf{X}, \Theta)]] , \end{aligned}$$

where we used a symmetry argument in the last equality. Observe now that

$$\begin{aligned} \mathbb{E}_{\Theta}^2 [W_{n1}(\mathbf{X}, \Theta)] &= \mathbb{E}_{\Theta} [W_{n1}(\mathbf{X}, \Theta)] \mathbb{E}_{\Theta'} [W_{n1}(\mathbf{X}, \Theta')] \\ &\quad \text{(where } \Theta' \text{ is distributed as, and independent of, } \Theta\text{)} \\ &= \mathbb{E}_{\Theta, \Theta'} [W_{n1}(\mathbf{X}, \Theta) W_{n1}(\mathbf{X}, \Theta')] \\ &= \mathbb{E}_{\Theta, \Theta'} \left[ \frac{\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta)]} \mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta')]} }{N_n(\mathbf{X}, \Theta) N_n(\mathbf{X}, \Theta')} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta)} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta')} \right] \\ &\leq \mathbb{E}_{\Theta, \Theta'} \left[ \frac{\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')]} }{N_n(\mathbf{X}, \Theta) N_n(\mathbf{X}, \Theta')} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta)} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta')} \right]. \end{aligned}$$

Consequently,

$$\mathbb{E} [\bar{r}_n(\mathbf{X}) - \tilde{r}_n(\mathbf{X})]^2 \leq n\sigma^2 \mathbb{E} \left[ \frac{\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')]} }{N_n(\mathbf{X}, \Theta) N_n(\mathbf{X}, \Theta')} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta)} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta')} \right].$$

Therefore

$$\begin{aligned} & \mathbb{E} [\bar{r}_n(\mathbf{X}) - \tilde{r}_n(\mathbf{X})]^2 \\ & \leq n\sigma^2 \mathbb{E} \left[ \frac{\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')]} }{\left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}\right) \left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta')]} \right)} \right] \\ & = n\sigma^2 \mathbb{E} \left[ \mathbb{E} \left[ \frac{\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')]} }{\left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}\right)} \right. \right. \\ & \quad \left. \left. \times \frac{1}{\left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta')]} \right)} \mid \mathbf{X}, \mathbf{X}_1, \Theta, \Theta' \right] \right] \\ & = n\sigma^2 \mathbb{E} \left[ \mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')]} \mathbb{E} \left[ \frac{1}{\left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}\right)} \right. \right. \\ & \quad \left. \left. \times \frac{1}{\left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta')]} \right)} \mid \mathbf{X}, \mathbf{X}_1, \Theta, \Theta' \right] \right] \\ & = n\sigma^2 \mathbb{E} \left[ \mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')]} \mathbb{E} \left[ \frac{1}{\left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}\right)} \right. \right. \\ & \quad \left. \left. \times \frac{1}{\left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta')]} \right)} \mid \mathbf{X}, \Theta, \Theta' \right] \right] \end{aligned}$$

by the independence of the random variables  $\mathbf{X}, \mathbf{X}_1, \dots, \mathbf{X}_n, \Theta, \Theta'$ . Using the Cauchy-Schwarz inequality, the above conditional expectation can be upper bounded by

$$\begin{aligned} & \mathbb{E}^{1/2} \left[ \frac{1}{\left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}\right)^2} \mid \mathbf{X}, \Theta \right] \\ & \quad \times \mathbb{E}^{1/2} \left[ \frac{1}{\left(1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta')]} \right)^2} \mid \mathbf{X}, \Theta' \right] \\ & \leq \frac{3 \times 2^{2\lceil \log_2 k_n \rceil}}{n^2} \\ & \quad \text{(by Fact 4.2 and technical Lemma 4.1)} \\ & \leq \frac{12k_n^2}{n^2}. \end{aligned}$$

It follows that

$$\begin{aligned}
\mathbb{E} [\bar{r}_n(\mathbf{X}) - \tilde{r}_n(\mathbf{X})]^2 &\leq \frac{12\sigma^2 k_n^2}{n} \mathbb{E} [\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')]}] \\
&= \frac{12\sigma^2 k_n^2}{n} \mathbb{E} [\mathbb{E}_{\mathbf{X}_1} [\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')]}]] \\
&= \frac{12\sigma^2 k_n^2}{n} \mathbb{E} [\mathbb{P}_{\mathbf{X}_1} (\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta'))]. \quad (4.1)
\end{aligned}$$

Next, using the fact that  $\mathbf{X}_1$  is uniformly distributed over  $[0, 1]^d$ , we may write

$$\begin{aligned}
\mathbb{P}_{\mathbf{X}_1} (\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')) &= \lambda (A_n(\mathbf{X}, \Theta) \cap A_n(\mathbf{X}, \Theta')) \\
&= \prod_{j=1}^d \lambda (A_{nj}(\mathbf{X}, \Theta) \cap A_{nj}(\mathbf{X}, \Theta')),
\end{aligned}$$

where

$$A_n(\mathbf{X}, \Theta) = \prod_{j=1}^d A_{nj}(\mathbf{X}, \Theta) \quad \text{and} \quad A_n(\mathbf{X}, \Theta') = \prod_{j=1}^d A_{nj}(\mathbf{X}, \Theta').$$

On the other hand, we know (Fact 4.1) that, for all  $j = 1, \dots, d$ ,

$$\lambda (A_{nj}(\mathbf{X}, \Theta)) \stackrel{\mathcal{D}}{=} 2^{-K_{nj}(\mathbf{X}, \Theta)},$$

where, conditionally on  $\mathbf{X}$ ,  $K_{nj}(\mathbf{X}, \Theta)$  has a binomial  $\mathcal{B}(\lceil \log_2 k_n \rceil, p_{nj})$  distribution and, similarly,

$$\lambda (A_{nj}(\mathbf{X}, \Theta')) \stackrel{\mathcal{D}}{=} 2^{-K'_{nj}(\mathbf{X}, \Theta')},$$

where, conditionally on  $\mathbf{X}$ ,  $K'_{nj}(\mathbf{X}, \Theta')$  is binomial  $\mathcal{B}(\lceil \log_2 k_n \rceil, p_{nj})$  and independent of  $K_{nj}(\mathbf{X}, \Theta)$ . In the rest of the proof, to lighten notation, we write  $K_{nj}$  and  $K'_{nj}$  instead of  $K_{nj}(\mathbf{X}, \Theta)$  and  $K'_{nj}(\mathbf{X}, \Theta')$ , respectively. Clearly,

$$\begin{aligned}
\lambda (A_{nj}(\mathbf{X}, \Theta) \cap A_{nj}(\mathbf{X}, \Theta')) &\leq 2^{-\max(K_{nj}, K'_{nj})} \\
&= 2^{-K'_{nj}} 2^{-(K_{nj} - K'_{nj})_+}
\end{aligned}$$

and, consequently,

$$\prod_{j=1}^d \lambda (A_{nj}(\mathbf{X}, \Theta) \cap A_{nj}(\mathbf{X}, \Theta')) \leq 2^{-\lceil \log_2 k_n \rceil} \prod_{j=1}^d 2^{-(K_{nj} - K'_{nj})_+}$$

(since, by Fact 4.1,  $\sum_{j=1}^d K_{nj} = \lceil \log_2 k_n \rceil$ ). Plugging this inequality into (4.1) and applying Hölder's inequality, we obtain

$$\begin{aligned} \mathbb{E} [\bar{r}_n(\mathbf{X}) - \tilde{r}_n(\mathbf{X})]^2 &\leq \frac{12\sigma^2 k_n}{n} \mathbb{E} \left[ \prod_{j=1}^d 2^{-(K_{nj}-K'_{nj})_+} \right] \\ &= \frac{12\sigma^2 k_n}{n} \mathbb{E} \left[ \mathbb{E} \left[ \prod_{j=1}^d 2^{-(K_{nj}-K'_{nj})_+} \mid \mathbf{X} \right] \right] \\ &\leq \frac{12\sigma^2 k_n}{n} \mathbb{E} \left[ \prod_{j=1}^d \mathbb{E}^{1/d} \left[ 2^{-d(K_{nj}-K'_{nj})_+} \mid \mathbf{X} \right] \right]. \end{aligned}$$

Each term in the product may be bounded by technical Proposition 4.1, and this leads to

$$\begin{aligned} \mathbb{E} [\bar{r}_n(\mathbf{X}) - \tilde{r}_n(\mathbf{X})]^2 &\leq \frac{288\sigma^2 k_n}{\pi n} \prod_{j=1}^d \min \left( 1, \left[ \frac{\pi}{16 \lceil \log_2 k_n \rceil p_{nj} (1 - p_{nj})} \right]^{1/2d} \right) \\ &\leq \frac{288\sigma^2 k_n}{\pi n} \prod_{j=1}^d \min \left( 1, \left[ \frac{\pi \log 2}{16 (\log k_n) p_{nj} (1 - p_{nj})} \right]^{1/2d} \right). \end{aligned}$$

Using the assumption on the form of the  $p_{nj}$ , we finally conclude that

$$\mathbb{E} [\bar{r}_n(\mathbf{X}) - \tilde{r}_n(\mathbf{X})]^2 \leq C \sigma^2 \left( \frac{S^2}{S-1} \right)^{S/2d} (1 + \xi_n) \frac{k_n}{n (\log k_n)^{S/2d}},$$

where

$$C = \frac{288}{\pi} \left( \frac{\pi \log 2}{16} \right)^{S/2d}$$

and

$$1 + \xi_n = \prod_{j \in \mathcal{S}} \left[ (1 + \xi_{nj})^{-1} \left( 1 - \frac{\xi_{nj}}{S-1} \right)^{-1} \right]^{1/2d}.$$

Clearly, the sequence  $(\xi_n)$ , which depends on the  $\{(\xi_{nj}) : j \in \mathcal{S}\}$  only, tends to 0 as  $n$  tends to infinity.

### 4.3 Proof of Proposition 2.2

We start with the decomposition

$$\begin{aligned}
& \mathbb{E} [\tilde{r}_n(\mathbf{X}) - r(\mathbf{X})]^2 \\
&= \mathbb{E} \left[ \sum_{i=1}^n \mathbb{E}_{\Theta} [W_{ni}(\mathbf{X}, \Theta)] (r(\mathbf{X}_i) - r(\mathbf{X})) \right. \\
&\quad \left. + \left( \sum_{i=1}^n \mathbb{E}_{\Theta} [W_{ni}(\mathbf{X}, \Theta)] - 1 \right) r(\mathbf{X}) \right]^2 \\
&= \mathbb{E} \left[ \mathbb{E}_{\Theta} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) + \left( \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) - 1 \right) r(\mathbf{X}) \right] \right]^2 \\
&\leq \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) + \left( \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) - 1 \right) r(\mathbf{X}) \right]^2 \\
&\quad \text{(by Jensen's inequality).}
\end{aligned}$$

Consequently,

$$\begin{aligned}
& \mathbb{E} [\tilde{r}_n(\mathbf{X}) - r(\mathbf{X})]^2 \\
&= \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) \right]^2 + \mathbb{E} [r(\mathbf{X}) \mathbf{1}_{\mathcal{E}_n^c(\mathbf{X}, \Theta)}]^2 \\
&\leq \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) \right]^2 + \left[ \sup_{\mathbf{x} \in [0,1]^d} r^2(\mathbf{x}) \right] \mathbb{P}(\mathcal{E}_n^c(\mathbf{X}, \Theta)).
\end{aligned} \tag{4.2}$$

Let us examine the first term on the right-hand side of (4.2). Observe that

$$\begin{aligned}
& \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) \right]^2 \\
&\leq \mathbb{E} \left[ \sum_{i=1}^n \sqrt{W_{ni}(\mathbf{X}, \Theta)} \sqrt{W_{ni}(\mathbf{X}, \Theta)} |r(\mathbf{X}_i) - r(\mathbf{X})| \right]^2 \\
&\leq \mathbb{E} \left[ \left( \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) \right) \left( \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X}))^2 \right) \right] \\
&\leq \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X}))^2 \right] \\
&\quad \text{(since the weights are subprobability weights).}
\end{aligned}$$

Thus, denoting by  $\|\mathbf{X}\|_{\mathcal{S}}$  the norm of  $\mathbf{X}$  evaluated over the components in  $\mathcal{S}$ , we obtain

$$\begin{aligned}
& \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) \right]^2 \\
& \leq \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r^*(\mathbf{X}_{i\mathcal{S}}) - r^*(\mathbf{X}_{\mathcal{S}}))^2 \right] \\
& \leq L^2 \sum_{i=1}^n \mathbb{E} [W_{ni}(\mathbf{X}, \Theta) \|\mathbf{X}_i - \mathbf{X}\|_{\mathcal{S}}^2] \\
& = nL^2 \mathbb{E} [W_{n1}(\mathbf{X}, \Theta) \|\mathbf{X}_1 - \mathbf{X}\|_{\mathcal{S}}^2] \\
& \quad (\text{by symmetry}).
\end{aligned}$$

But

$$\begin{aligned}
& \mathbb{E} [W_{n1}(\mathbf{X}, \Theta) \|\mathbf{X}_1 - \mathbf{X}\|_{\mathcal{S}}^2] \\
& = \mathbb{E} \left[ \|\mathbf{X}_1 - \mathbf{X}\|_{\mathcal{S}}^2 \frac{\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta)]}}{N_n(\mathbf{X}, \Theta)} \mathbf{1}_{\mathcal{E}_n(\mathbf{X}, \Theta)} \right] \\
& = \mathbb{E} \left[ \|\mathbf{X}_1 - \mathbf{X}\|_{\mathcal{S}}^2 \frac{\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta)]}}{1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}} \right] \\
& = \mathbb{E} \left[ \mathbb{E} \left[ \|\mathbf{X}_1 - \mathbf{X}\|_{\mathcal{S}}^2 \frac{\mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta)]}}{1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}} \mid \mathbf{X}, \mathbf{X}_1, \Theta \right] \right] \\
& = \mathbb{E} \left[ \|\mathbf{X}_1 - \mathbf{X}\|_{\mathcal{S}}^2 \mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta)]} \mathbb{E} \left[ \frac{1}{1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}} \mid \mathbf{X}, \mathbf{X}_1, \Theta \right] \right] \\
& = \mathbb{E} \left[ \|\mathbf{X}_1 - \mathbf{X}\|_{\mathcal{S}}^2 \mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta)]} \mathbb{E} \left[ \frac{1}{1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}} \mid \mathbf{X}, \Theta \right] \right] \\
& \quad (\text{by the independence of the random variables } \mathbf{X}, \mathbf{X}_1, \dots, \mathbf{X}_n, \Theta).
\end{aligned}$$

By Fact 4.2 and technical Lemma 4.1,

$$\mathbb{E} \left[ \frac{1}{1 + \sum_{i=2}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{X}, \Theta)]}} \mid \mathbf{X}, \Theta \right] \leq \frac{2^{\lceil \log_2 k_n \rceil}}{n} \leq \frac{2k_n}{n}.$$

Consequently,

$$\begin{aligned}
& \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) \right]^2 \\
& \leq 2L^2 k_n \mathbb{E} [\|\mathbf{X}_1 - \mathbf{X}\|_{\mathcal{S}}^2 \mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta)]}].
\end{aligned}$$

Letting

$$A_n(\mathbf{X}, \Theta) = \prod_{j=1}^d A_{nj}(\mathbf{X}, \Theta),$$

we obtain

$$\begin{aligned} & \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) \right]^2 \\ & \leq 2L^2 k_n \sum_{j \in \mathcal{S}} \mathbb{E} \left[ |\mathbf{X}_1^{(j)} - \mathbf{X}^{(j)}|^2 \mathbf{1}_{[\mathbf{X}_1 \in A_n(\mathbf{X}, \Theta)]} \right] \\ & = 2L^2 k_n \sum_{j \in \mathcal{S}} \mathbb{E} \left[ \rho_j(\mathbf{X}, \mathbf{X}_1, \Theta) \mathbb{E}_{\mathbf{X}_1^{(j)}} \left[ |\mathbf{X}_1^{(j)} - \mathbf{X}^{(j)}|^2 \mathbf{1}_{[\mathbf{X}_1^{(j)} \in A_{nj}(\mathbf{X}, \Theta)]} \right] \right] \end{aligned}$$

where, in the last equality, we set

$$\rho_j(\mathbf{X}, \mathbf{X}_1, \Theta) = \prod_{t=1, \dots, d, t \neq j} \mathbf{1}_{[\mathbf{X}_1^{(t)} \in A_{nt}(\mathbf{X}, \Theta)]}.$$

Therefore, using the fact that  $\mathbf{X}_1$  is uniformly distributed over  $[0, 1]^d$ ,

$$\begin{aligned} & \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) \right]^2 \\ & \leq 2L^2 k_n \sum_{j \in \mathcal{S}} \mathbb{E} \left[ \rho_j(\mathbf{X}, \mathbf{X}_1, \Theta) \lambda^3(A_{nj}(\mathbf{X}, \Theta)) \right]. \end{aligned}$$

Observing that

$$\begin{aligned} & \lambda(A_{nj}(\mathbf{X}, \Theta)) \times \mathbb{E}_{[\mathbf{X}_1^{(t)} : t=1, \dots, d, t \neq j]} [\rho_j(\mathbf{X}, \mathbf{X}_1, \Theta)] \\ & = \lambda(A_n(\mathbf{X}, \Theta)) \\ & = 2^{-\lceil \log_2 k_n \rceil} \\ & \quad (\text{Fact 4.2}), \end{aligned}$$

we are led to

$$\begin{aligned} & \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) \right]^2 \\ & \leq 2L^2 \sum_{j \in \mathcal{S}} \mathbb{E} \left[ \lambda^2(A_{nj}(\mathbf{X}, \Theta)) \right] \\ & = 2L^2 \sum_{j \in \mathcal{S}} \mathbb{E} \left[ 2^{-2K_{nj}(\mathbf{X}, \Theta)} \right] \\ & = 2L^2 \sum_{j \in \mathcal{S}} \mathbb{E} \left[ \mathbb{E} \left[ 2^{-2K_{nj}(\mathbf{X}, \Theta)} \mid \mathbf{X} \right] \right], \end{aligned}$$

where, conditionally on  $\mathbf{X}$ ,  $K_{nj}(\mathbf{X}, \Theta)$  has a binomial  $\mathcal{B}(\lceil \log_2 k_n \rceil, p_{nj})$  distribution (Fact 4.1). Consequently,

$$\begin{aligned}
& \mathbb{E} \left[ \sum_{i=1}^n W_{ni}(\mathbf{X}, \Theta) (r(\mathbf{X}_i) - r(\mathbf{X})) \right]^2 \\
& \leq 2L^2 \sum_{j \in \mathcal{S}} (1 - 0.75p_{nj})^{\lceil \log_2 k_n \rceil} \\
& \leq 2L^2 \sum_{j \in \mathcal{S}} \exp \left( -\frac{0.75}{\log 2} p_{nj} \log k_n \right) \\
& = 2L^2 \sum_{j \in \mathcal{S}} \frac{1}{k_n^{\frac{0.75}{S \log 2} (1 + \xi_{nj})}} \\
& \leq \frac{2SL^2}{k_n^{\frac{0.75}{S \log 2} (1 + \gamma_n)}},
\end{aligned}$$

with  $\gamma_n = \min_{j \in \mathcal{S}} \xi_{nj}$ .

To finish the proof, it remains to bound the second term on the right-hand side of (4.2), which is easier. Just note that

$$\begin{aligned}
\mathbb{P}(\mathcal{E}_n^c(\mathbf{X}, \Theta)) &= \mathbb{P} \left( \sum_{i=1}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{x}, \Theta)]} = 0 \right) \\
&= \mathbb{E} \left[ \mathbb{P} \left( \sum_{i=1}^n \mathbf{1}_{[\mathbf{X}_i \in A_n(\mathbf{x}, \Theta)]} = 0 \mid \mathbf{X}, \Theta \right) \right] \\
&= (1 - 2^{-\lceil \log_2 k_n \rceil})^n \\
&\quad \text{(by Fact 4.2)} \\
&\leq e^{-n/2k_n}.
\end{aligned}$$

Putting all the pieces together, we finally conclude that

$$\mathbb{E} [\tilde{r}_n(\mathbf{X}) - r(\mathbf{X})]^2 \leq \frac{2SL^2}{k_n^{\frac{0.75}{S \log 2} (1 + \gamma_n)}} + \left[ \sup_{\mathbf{x} \in [0,1]^d} r^2(\mathbf{x}) \right] e^{-n/2k_n},$$

as desired.

#### 4.4 Some technical results

The following result is an extension of Lemma 4.1 in Györfi et al. [22]. Its proof is given here for the sake of completeness.

**Lemma 4.1** *Let  $Z$  be a binomial  $\mathcal{B}(N, p)$  random variable, with  $p \in (0, 1]$ . Then*

(i)

$$\mathbb{E} \left[ \frac{1}{1+Z} \right] \leq \frac{1}{(N+1)p}.$$

(ii)

$$\mathbb{E} \left[ \frac{1}{Z} \mathbf{1}_{[Z \geq 1]} \right] \leq \frac{2}{(N+1)p}.$$

(iii)

$$\mathbb{E} \left[ \frac{1}{1+Z^2} \right] \leq \frac{3}{(N+1)(N+2)p^2}.$$

**Proof of Lemma 4.1** To prove statement (i), we write

$$\begin{aligned} \mathbb{E} \left[ \frac{1}{1+Z} \right] &= \sum_{j=0}^N \frac{1}{1+j} \binom{N}{j} p^j (1-p)^{N-j} \\ &= \frac{1}{(N+1)p} \sum_{j=0}^N \binom{N+1}{j+1} p^{j+1} (1-p)^{N-j} \\ &\leq \frac{1}{(N+1)p} \sum_{j=0}^{N+1} \binom{N+1}{j} p^j (1-p)^{N+1-j} \\ &= \frac{1}{(N+1)p}. \end{aligned}$$

The second statement follows from the inequality

$$\mathbb{E} \left[ \frac{1}{Z} \mathbf{1}_{[Z \geq 1]} \right] \leq \mathbb{E} \left[ \frac{2}{1+Z} \right]$$

and the third one by observing that

$$\mathbb{E} \left[ \frac{1}{1+Z^2} \right] = \sum_{j=0}^N \frac{1}{1+j^2} \binom{N}{j} p^j (1-p)^{N-j}.$$

Therefore

$$\begin{aligned}
\mathbb{E} \left[ \frac{1}{1+Z^2} \right] &= \frac{1}{(N+1)p} \sum_{j=0}^N \frac{1+j}{1+j^2} \binom{N+1}{j+1} p^{j+1} (1-p)^{N-j} \\
&\leq \frac{3}{(N+1)p} \sum_{j=0}^N \frac{1}{2+j} \binom{N+1}{j+1} p^{j+1} (1-p)^{N-j} \\
&\leq \frac{3}{(N+1)p} \sum_{j=0}^{N+1} \frac{1}{1+j} \binom{N+1}{j} p^j (1-p)^{N+1-j} \\
&\leq \frac{3}{(N+1)(N+2)p^2} \\
&\quad \text{(by (i)).}
\end{aligned}$$

■

**Lemma 4.2** *Let  $Z_1$  and  $Z_2$  be two independent binomial  $\mathcal{B}(N, p)$  random variables. Set, for all  $z \in \mathbb{C}^*$ ,  $\varphi(z) = \mathbb{E}[z^{Z_1 - Z_2}]$ .*

(i) *For all  $z \in \mathbb{C}^*$ ,*

$$\varphi(z) = [p(1-p)(z + z^{-1}) + 1 - 2p(1-p)]^N.$$

(ii) *For all  $j \in \mathbb{N}$ ,*

$$\mathbb{P}(Z_1 - Z_2 = j) = \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi(z)}{z^{j+1}} dz,$$

*where  $\Gamma$  is the positively oriented unit circle.*

(iii) *For all  $d \geq 1$ ,*

$$\mathbb{E} [2^{-d(Z_1 - Z_2)_+}] \leq \frac{24}{\pi} \int_0^1 \exp(-4Np(1-p)t^2) dt.$$

**Proof of Lemma 4.2** Statement (i) is clear and (ii) is an immediate consequence of Cauchy's integral formula (Rudin [29]). To prove statement

(iii), write

$$\begin{aligned}
\mathbb{E} [2^{-d(Z_1 - Z_2)_+}] &= \sum_{j=0}^N 2^{-dj} \mathbb{P}((Z_1 - Z_2)_+ = j) \\
&= \sum_{j=0}^N 2^{-dj} \mathbb{P}(Z_1 - Z_2 = j) \\
&\leq \sum_{j=0}^{\infty} 2^{-dj} \mathbb{P}(Z_1 - Z_2 = j) \\
&= \frac{1}{2\pi i} \int_{\Gamma} \frac{\varphi(z)}{z} \sum_{j=0}^{\infty} \left(\frac{2^{-d}}{z}\right)^j dz \\
&\quad \text{(by statement (ii))} \\
&= \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{\varphi(e^{i\theta})}{1 - 2^{-d}e^{-i\theta}} d\theta \\
&\quad \text{(by setting } z = e^{i\theta}, \theta \in [-\pi, \pi]) \\
&= \frac{2^{d-1}}{\pi} \int_{-\pi}^{\pi} [1 + 2p(1-p)(\cos \theta - 1)]^N \frac{e^{i\theta}}{2^d e^{i\theta} - 1} d\theta \\
&\quad \text{(by statement (i)).}
\end{aligned}$$

Noting that

$$\frac{e^{i\theta}}{2^d e^{i\theta} - 1} = \frac{2^d - e^{i\theta}}{2^{2d} - 2^{d+1} \cos \theta + 1},$$

we obtain

$$\begin{aligned}
\mathbb{E} [2^{-d(Z_1 - Z_2)_+}] \\
\leq \frac{2^{d-1}}{\pi} \int_{-\pi}^{\pi} [1 + 2p(1-p)(\cos \theta - 1)]^N \frac{2^d - \cos \theta}{2^{2d} - 2^{d+1} \cos \theta + 1} d\theta.
\end{aligned}$$

The bound

$$\frac{2^d - \cos \theta}{2^{2d} - 2^{d+1} \cos \theta + 1} \leq \frac{2^d + 1}{(2^d - 1)^2}$$

leads to

$$\begin{aligned}
& \mathbb{E} [2^{-d(Z_1 - Z_2)_+}] \\
& \leq \frac{2^{d-1}(2^d + 1)}{\pi(2^d - 1)^2} \int_{-\pi}^{\pi} [1 + 2p(1 - p)(\cos \theta - 1)]^N d\theta \\
& = \frac{2^d(2^d + 1)}{\pi(2^d - 1)^2} \int_0^{\pi} [1 + 2p(1 - p)(\cos \theta - 1)]^N d\theta \\
& = \frac{2^d(2^d + 1)}{\pi(2^d - 1)^2} \int_0^{\pi} [1 - 4p(1 - p) \sin^2(\theta/2)]^N d\theta \\
& \quad (\cos \theta - 1 = -2 \sin^2(\theta/2)) \\
& = \frac{2^{d+1}(2^d + 1)}{\pi(2^d - 1)^2} \int_0^{\pi/2} [1 - 4p(1 - p) \sin^2 \theta]^N d\theta.
\end{aligned}$$

Using the elementary inequality  $(1 - z)^N \leq e^{-Nz}$  for  $z \in [0, 1]$  and the change of variable

$$t = \tan(\theta/2),$$

we finally obtain

$$\begin{aligned}
& \mathbb{E} [2^{-d(Z_1 - Z_2)_+}] \\
& \leq \frac{2^{d+2}(2^d + 1)}{\pi(2^d - 1)^2} \int_0^1 \exp\left(-\frac{16Np(1 - p)t^2}{(1 + t^2)^2}\right) \frac{1}{1 + t^2} dt \\
& \leq C_d \int_0^1 \exp(-4Np(1 - p)t^2) dt,
\end{aligned}$$

with

$$C_d = \frac{2^{d+2}(2^d + 1)}{\pi(2^d - 1)^2}.$$

The conclusion follows by observing that  $C_d \leq 24/\pi$  for all  $d \geq 1$ . ■

Evaluating the integral in statement (iii) of Lemma 4.2 leads to the following proposition:

**Proposition 4.1** *Let  $Z_1$  and  $Z_2$  be two independent binomial  $\mathcal{B}(N, p)$  random variables, with  $p \in (0, 1)$ . Then, for all  $d \geq 1$ ,*

$$\mathbb{E} [2^{-d(Z_1 - Z_2)_+}] \leq \frac{24}{\pi} \min\left(1, \sqrt{\frac{\pi}{16Np(1 - p)}}\right).$$

## References

- [1] D. Amaratunga, J. Cabrera, and Y.S. Lee. Enriched random forests. *Bioinformatics*, 24:2010–2014, 2008.
- [2] Y. Amit and D. Geman. Shape quantization and recognition with randomized trees. *Neural Computation*, 9:1545–1588, 1997.
- [3] G. Biau, F. Cérou, and A. Guyader. On the rate of convergence of the bagged nearest neighbor estimate. *Journal of Machine Learning Research*, 11:687–712, 2010.
- [4] G. Biau and L. Devroye. *On the layered nearest neighbour estimate, the bagged nearest neighbour estimate and the random forest method in regression and classification*. Technical Report, Université Pierre et Marie Curie, Paris, 2008.
- [5] G. Biau, L. Devroye, and G. Lugosi. Consistency of random forests and other averaging classifiers. *Journal of Machine Learning Research*, 9:2015–2033, 2008.
- [6] P. Bickel, Y. Ritov, and A. Tsybakov. Simultaneous analysis of Lasso and Dantzig selector. *The Annals of Statistics*, 37:1705–1732, 2009.
- [7] L. Breiman. Bagging predictors. *Machine Learning*, 24:123–140, 1996.
- [8] L. Breiman. *Some infinity theory for predictor ensembles*. Technical Report 577, UC Berkeley, 2000.
- [9] L. Breiman. Random forests. *Machine Learning*, 45:5–32, 2001.
- [10] L. Breiman. *Consistency for a simple model of random forests*. Technical Report 670, UC Berkeley, 2004.
- [11] L. Breiman, J.H. Friedman, R.A. Olshen, and C.J. Stone. *Classification and Regression Trees*. Wadsworth Statistics/Probability Series, Belmont, 1984.
- [12] A.M. Bruckstein, D.L. Donoho, and M. Elad. From sparse solutions of systems of equations to sparse modeling of signals and images. *SIAM Review*, 51:34–81, 2009.
- [13] P. Bühlmann and B. Yu. Analyzing bagging. *The Annals of Statistics*, 30:927–961, 2002.

- [14] A. Buja and W. Stuetzle. Observations on bagging. *Statistica Sinica*, 16:323–352, 2006.
- [15] F. Bunea, A. Tsybakov, and M. Wegkamp. Sparsity oracle inequalities for the Lasso. *Electronic Journal of Statistics*, 1:169–194, 2007.
- [16] E.J. Candès and T. Tao. The Dantzig selector: Statistical estimation when  $p$  is much larger than  $n$ . *The Annals of Statistics*, 35:2313–2351, 2005.
- [17] L. Devroye, L. Györfi, and G. Lugosi. *A Probabilistic Theory of Pattern Recognition*. Springer-Verlag, New York, 1996.
- [18] R. Diaz-Uriarte and S.A. de Andrés. Gene selection and classification of microarray data using random forest. *BMC Bioinformatics*, 7:1471–2105, 2006.
- [19] T.G. Dietterich. An experimental comparison of three methods for constructing ensembles of decision trees: bagging, boosting, and randomization. *Machine Learning*, 40:139–157, 2000.
- [20] Y. Freund and R. Shapire. Experiments with a new boosting algorithm. In L. Saitta, editor, *Machine Learning: Proceedings of the 13th International Conference*, pages 148–156, San Francisco, 1996. Morgan Kaufmann.
- [21] R. Genuer, J.-M. Poggi, and C. Tuleau. *Random Forests: Some methodological insights*. Research Report 6729, INRIA Saclay-Ile-de-France, 2008.
- [22] L. Györfi, M. Kohler, A. Krzyżak, and H. Walk. *A Distribution-Free Theory of Nonparametric Regression*. Springer-Verlag, New York, 2002.
- [23] T.K. Ho. The random subspace method for constructing decision forests. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 20:832–844, 1998.
- [24] I.A. Ibragimov and R.Z. Khasminskii. On nonparametric estimation of regression. *Doklady Akademii Nauk SSSR*, 252:780–784, 1980.
- [25] I.A. Ibragimov and R.Z. Khasminskii. *Statistical Estimation: Asymptotic Theory*. Springer-Verlag, New York, 1981.

- [26] I.A. Ibragimov and R.Z. Khasminskii. On the bounds for quality of nonparametric regression function estimation. *Theory of Probability and its Applications*, 27:81–94, 1982.
- [27] Y. Lin and Y. Jeon. Random forests and adaptive nearest neighbors. *Journal of the American Statistical Association*, 101:578–590, 2006.
- [28] N. Meinshausen. Quantile regression forests. *Journal of Machine Learning Research*, 7:983–999, 2006.
- [29] W. Rudin. *Real and Complex Analysis, 3rd Edition*. McGraw-Hill, New York, 1987.
- [30] J. Shawe-Taylor and N. Cristianini. *Kernel Methods for Pattern Analysis*. Cambridge University Press, Cambridge, 2004.
- [31] V. Svetnik, A. Liaw, C. Tong, J. Culberson, R. Sheridan, and B. Feuston. Random forest: A classification and regression tool for compound classification and QSAR modeling. *Journal of Chemical Information and Computer Sciences*, 43:1947–1958, 2003.
- [32] R. Tibshirani. Regression shrinkage and selection via the Lasso. *Journal of the Royal Statistical Society, Series B*, 58:267–288, 1996.
- [33] L.J. van't Veer, H. Dai, M.J. van de Vijver, Y.D. He, A.A.M. Hart, M. Mao, H.L. Peterse, K. van der Kooy, M.J. Marton, A.T. Witteveen, G.J. Schreiber, R.M. Kerkhoven, C. Roberts, P.S. Linsley, R. Bernards, and S.H. Friend. Gene expression profiling predicts clinical outcome of breast cancer. *Nature*, 415:530–536, 2002.